

С. Баранитаран, Л. Чандракумар
**МОРФОЛОГИЧЕСКИЕ ОСОБЕННОСТИ ОТВЕРСТИЙ И УЗЕЛКОВ
ПОЛУЛУННЫХ ЗАСЛОНОК КЛАПАНА АОРТЫ ЧЕЛОВЕКА**

Научный руководитель: канд. мед. наук, доц. А.А. Пасюк

Кафедра нормальной анатомии

Белорусский государственный медицинский университет, г. Минск

S. Baraneetharan, L. Chandrakumar
**MORPHOLOGICAL FEATURES OF FENESTRA AND NODULES
ON THE AORTIC VALVE LEAFLETS OF HUMANS**

Tutor: PhD associate professor H.A. Pasiuk

Department of Normal Anatomy

Belarusian State Medical University, Minsk

Резюме. В статье представлены результаты морфологического исследования особенностей узелков и отверстий полулунных заслонок клапана аорты на 38 аутопсийных сердцах взрослого человека. Установлено положение узелка на полулунной заслонке, выявлены основные формы узелков и определена частота их встречаемости. Представлены сведения о частоте встречаемости и закономерностях распределения отверстий полулунных заслонок клапана аорты.

Ключевые слова: клапан аорты, полулунная заслонка, узелок полулунной заслонки, отверстие полулунной заслонки, человек, морфология.

Resume. The article presents the results of a morphological study on the characteristics of nodules and fenestrae in the semilunar cusps of the aortic valve, based on 38 autopsied adult human hearts. The study determines the position of the nodule on the semilunar cusp, identifies the main nodule shapes, and establishes their occurrence frequency. Data on the prevalence and distribution patterns of fenestrae in the semilunar cusps of the aortic valve are also provided.

Keywords: aortic valve, semilunar leaflet, nodules of semilunar leaflet, fenestra of semilunar leaflet, human, morphology.

Relevance. The aortic root serves as the anatomical connection between the left ventricle and the ascending aorta, playing a crucial role in cardiovascular function. It consists of the aortic valve leaflets, which are supported by the aortic sinuses (Valsalva), and the interleaflet triangles that lie between their basal attachments, ensuring structural integrity and optimal blood flow dynamics [1]. Nodules of semilunar leaflet of aortic valve are the fibrous thickenings located at the midpoints of the cusp. They ensure complete valve closure and, together with the semilunar lunulas, form the coaptation area to prevent backflow. Size and integrity variations can indicate underlying valve disease, with enlarged or calcified nodules leading to aortic regurgitation [2]. Fenestrae are small cusp perforations, are normal in small numbers; however, excessive size or quantity weakens cusps, potentially causing regurgitation or prolapse [3]. Increased fenestrae near commissures raise tear risks and insufficiency, highlighting the importance of fenestrae characteristics in maintaining aortic valve function and predicting disease [4]. This study examines the structural characteristics of semilunar leaflets, nodules, and fenestrae of semilunar leaflet of aortic valve, aiming to refine diagnostic accuracy and support the development of targeted therapeutic strategies in cardiology.

Aim: to establish the range of anatomical variations in aortic valve nodules and fenestrae by characterizing their variant anatomy, nodule location, and frequency of fenestrae occurrence.

Objectives:

1. Determine the patterns of location of the nodules of semilunar leaflet of the aortic valve.
2. Determine the dominant forms of semilunar leaflet nodules.
3. Study the frequency of occurrence of fenestrae of semilunar leaflet of the aortic valve.

Material and methods. Morphological and morphometric methods were used to evaluate the characteristics of aortic valves in 38 post-mortem human hearts (68,50 [52,00; 78,50] years old). Measurements included lunula lengths (left and right), nodule width, and nodule height. Data were compiled and analysed using Microsoft Excel and Statistica 10.0.

Results and their discussion. The study found that the median nodule width across the aortic valve leaflets ranged from 2.00 to 4.00 mm, with notable variations among different leaflets. In terms of location, nodules exhibited distinct positional tendencies depending on the leaflet type. In the left coronary leaflet, nodules were predominantly shifted left in 43.75% of cases. In contrast, nodules in the non-coronary leaflet were more frequently shifted right (45.45% of cases), while those in the right coronary leaflet were centrally positioned in 37.84% of cases. These findings challenge the traditional assumption that nodules are always midpoint-centered, providing new perspectives on anatomical variability in aortic valve structures.




Fig. 1 – Analysis of Prevalence of location of Nodules




Fig. 2 – Analysis of prevalence of nodules <1mm (%)

The study identified five distinct nodule shapes within the aortic valve leaflets: brush/papillary, round, linear, small protrusion, and triangular. Notably, linear nodules were absent in the left coronary leaflet, while they were present in both the right coronary and non-coronary leaflets, suggesting structural variability across leaflet types. Among all observed shapes, brush/papillary and round nodules emerged as the most prevalent, appearing more frequently than other formations across all three leaflets. These findings highlight the diverse morphological characteristics of aortic valve nodules, contributing to a more refined understanding of their structural differences and potential clinical implications.




Fig. 3 – Analysis of Shape of Nodule (%)




Fig. 4 – Shapes of the nodules of semilunar leaflets in the sample

The study revealed notable variability in lunula length across the aortic valve leaflets. The left lunula length demonstrated a range of 11.00–15.00 mm in the left coronary leaflet, 12.00–16.00 mm in the right coronary leaflet, and 12.00–16.00 mm in the non-coronary leaflet, indicating slight differences in structural adaptation across leaflets. The right lunula length, however, was generally greater, spanning 12.00–17.00 mm across all leaflets, suggesting an anatomical tendency for increased length on the right side. These findings emphasize the structural diversity of aortic valve components.

Moreover, the study found that fenestrae, or small perforations in aortic valve leaflets, were present in 26.32% of the examined valves, highlighting their structural variability. The left coronary leaflet exhibited the highest incidence of fenestrae at 13.16%, followed by the right coronary leaflet at 10.53%, and the non-coronary leaflet at 4.00%, indicating an unequal distribution across leaflets. Among the cases with fenestrae, 50% showed perforations in only one leaflet, while 20% exhibited fenestrae in two leaflets, and 10% displayed perforations across all three leaflets. These findings suggest that fenestrae occur naturally, but their distribution and prevalence could have implications for valve integrity and function, particularly in relation to structural weakening and potential regurgitation risks.




Fig. 5 – Analysis of distribution and incidence of the fenestra of semilunar leaflets of aortic valve

The study underscores the importance of morphological variability in aortic valve nodules, challenging the traditional perspective that nodules are exclusively midpoint centered. The observed positional shifts suggest structural adaptability, which may influence valve closure efficiency. Additionally, the presence of increased fenestrae near commissures raises concerns about potential tear risks, as these perforations can weaken leaflet integrity over time, possibly leading to aortic regurgitation.

Conclusion:

1. Nodule placement varies across leaflets – Nodules tend to shift left in the left coronary leaflet, right in the non-coronary leaflet, and remain centrally located in the right coronary leaflet, challenging the traditional midpoint-only assumption.

2. Brush/papillary and round nodules are most common – These two shapes predominate across all leaflets, while linear nodules are absent in the left coronary leaflet, suggesting structural specialization.

3. Fenestrae distribution is uneven – Left coronary leaflets exhibit the highest frequency of fenestrae, followed by right coronary and non-coronary leaflets, emphasizing structural vulnerabilities and potential clinical implications.

Literature

1. Friedman B., Hathaway B. M. Fenestration of the semilunar cusps, and “functional” aortic and pulmonic valve insufficiency //The American Journal of Medicine. – 1958. – Т. 24. – №. 4. – С. 549-558.

2. Katsi V. et al. Aortic valve: anatomy and structure and the role of vasculature in the degenerative process //Acta Cardiologica. – 2021. – Т. 76. – №. 4. – С. 335-348.

3. Matthews R. J., Darvill JR F. T. Fenestrations of the aortic valve cusps as a cause of aortic insufficiency and spontaneous aortic valve cusp rupture //Annals of internal medicine. – 1956. – Т. 44. – №. 5. – С. 993-1002.

4. Tatar C. M., Ispas M. Anatomy and function of normal aortic valvular complex //Calcific aortic valve disease. – 2013. – С. 31.