МИНИСТЕРСТВО ЗДРАВООХРАНЕНИЯ РЕСПУБЛИКИ БЕЛАРУСЬ БЕЛОРУССКИЙ ГОСУДАРСТВЕННЫЙ МЕДИЦИНСКИЙ УНИВЕРСИТЕТ кафедра общей гигиены

ОБЩАЯ ГИГИЕНА

GENERAL HYGIENE

Лабораторный практикум

2-е издание, исправленное и дополненное



Минск БГМУ 2024

Рекомендовано Научно-методическим советом университета в качестве лабораторного практикума 21.03.2024 г., протокол № 15

А в т о р ы: ст. преп. К. В. Богданович; ст. преп. Е. И. Цимберова; ст. преп. А. Ю. Полещук; канд. мед. наук О. Н. Петровская; канд. мед. наук Я. Н. Борисевич; канд. мед. наук Н. В. Рябова

Рецензенты: канд. мед. наук, доц., зав. каф. гигиены детей и подростков Т. С. Борисова; канд. мед. наук, доц., зав. каф. гигиены труда И. П. Семёнов; каф. гигиены труда

Общая гигиена = General hygiene : лабораторный практикум / К. В. Богдано-О-28 вич [и др.]. – 2-е изд., испр. и доп. – Минск : БГМУ, 2024. – 108 с.

ISBN 978-985-21-1649-7.

Содержит информацию для подготовки к практическим занятиям по дисциплине «Общая гигиена». Представлены алгоритмы, справочные сведения, методики выполнения практических работ по основным темам дисциплины. Включены вопросы и задания для подготовки к коллоквиумам и итоговым занятиям. Первое издание вышло в 2023 году.

Предназначен для студентов 2-го курса медицинского факультета иностранных учащихся, изучающих общую гигиену на английском языке.

УДК 613(076.5)(075.8)-054.6 ББК 51.20я73

Учебное издание

Богданович Ксения Викторовна Цимберова Екатерина Игоревна Полещук Александра Юрьевна и др.

ОБЩАЯ ГИГИЕНА

GENERAL HYGIENE

Лабораторный практикум

На английском языке

2-е издание, исправленное и дополненное

Ответственная за выпуск Н. Л. Бацукова Переводчик К. В. Богданович Компьютерная вёрстка А. В. Янушкевич

Подписано в печать 11.09.24. Формат 60×84/8. Бумага писчая «Снегурочка». Ризография. Гарнитура «Times».

Усл. печ. л. 12,55. Уч.-изд. л. 4,1. Тираж 229 экз. Заказ 481.

Издатель и полиграфическое исполнение: учреждение образования «Белорусский государственный медицинский университет». Свидетельство о государственной регистрации издателя, изготовителя, распространителя печатных изданий № 1/187 от 24.11.2023.

Ул. Ленинградская, 6, 220006, Минск.

ISBN 978-985-21-1649-7

© УО «Белорусский государственный медицинский университет», 2024

PHYSICAL PROPERTIES OF ATMOSPHERIC AIR. RESEARCH METHODS AND HYGIENIC ASSESSMENT OF MICROCLIMATE PARAMETERS IN RESIDENTIAL PREMISES AND MEDICAL INSTITUTIONS (TEMPERATURE, HUMIDITY)

Learning objectives:

Date_____

1. To learn definition and features of weather and climate, climate classification.
developing during the acclimatization.
3. To learn definition of «microclimate» and factors, which influence its formation.
4. To learn basics of human heat exchange and thermoregulation.
5. To master methods of determination and hygienic assessment of temperature and humidity.
THEORETICAL SECTION
Give a definition
Climate —
Weather —
Microclimate
Acclimatization —
Absolute humidity —
Maximal humidity —
Relative humidity
Saturation deficiency —

PRACTICAL WORK

The microclimate of premises and its hygienic estimation.

1. Definition of a Temperature Mode.

The air temperature is measured by _____

Standard (normal) values = _____

To define a temperature mode of premises it is necessary to take air temperature in the following points: in the center and at the external and internal walls at the distance of 10 cm from them and at the level of 0.1-1-1.5 m from the floor. The obtained data are written to the protocol and analyzed on vertical and horizontal lines.

Actual values = _____

Conclusion:

2. Hygienic assessment of humidity.

The humidity is measured by

Standard (normal) values = _____

a) Finding humidity using table.

To assess humidity, use «Table for determining air humidity according to August psychrometer readings».

Actual value (relative humidity by the table) = _____

b) Finding humidity using devices.

Temperature of the dry thermometer = _____ Temperature of the wet thermometer = _____

1. K (absolute humidity) = $f - a \times (t_1 - t_2) \times B$

f — the maximum humidity (temperature of the wet thermometer); a — psychometric factor 0.0011; t₁ — temperature of the dry thermometer; t₂ — temperature of the wet thermometer; B — barometric pressure in mm Hg

Actual value (absolute humidity) = _____

2. R (relative humidity) = $K/F \times 100 \%$

K — absolute humidity; F — the maximum humidity (temperature of the dry thermometer) Actual value (relative humidity) = _____

3. D_{physic} (physical deficiency of saturation) = $F_{t room} - K$ Actual value (deficiency of saturation) = _____ 4. Dew-point temperature (equate the obtained value of absolute humidity to maximum and find the required temperature in the table)

Actual value (dew-point temperature) = _____

Conclusion:			
-			

PROBLEMS SOLVING

Example. Indications of dry and wet bulbs of aspirating August psychrometer in the living room are +19 °C and +18 °C respectively. Atmospheric pressure is 740 mmHg.

Determine the absolute humidity, relative humidity, saturation deficiency, dew point temperature. Compare the result with the hygienic standard.

F (the maximum humidity of the dry thermometer) = 16.48 mmHg (table) f (the maximum humidity of the wet thermometer) = 15.48 mmHg (table)

1. $K = f - a \times (t_1 - t_2) \times B = 15.48 - (0.0011 \times (19 - 18) \times 740) = 14.67 \text{ mmHg}$

2. $R = K/F \times 100 \% = 14.67/16.48 \times 100 \% = 90 \%$

- 3. $D_{physic} = F_{t room} K = 16.48 14.67 = 1.814 \text{ mmHg}$
- 4. Dew-point temperature = 17.2 °C

Conclusion: During the hygienic assessment of the microclimate parameters, it was found that the relative humidity exceeds the norm (actual = 90 %, normal = < 75 %).

Tasks. 1. Indications of dry and wet bulbs of aspirating August psychrometer in the classroom are +22 °C and +18 °C respectively. Atmospheric pressure is 755 mmHg.

Determine the relative humidity (according to the formula and table), saturation deficiency, asses them. What is the dew point temperature?

Perform hygienic assessment of calculated data (compare the result with the hygienic standard).

2. The parameters of microclimate in two closed rooms are: in the first — the air temperature is 22 °C, and the relative humidity — 61 %, in the second — 16 °C and 78 %, respectively. In which of the rooms is the value of physical saturation deficiency higher?

KEY QUESTIONS

1. Weather and climate, features and definition. The notion of climatic factors. Climate classification, their hygienic characteristics.

2. The mechanism of thermoregulation in the human organism and ways of heat transfer. Hygienic significance of the atmospheric and indoor air, its role in the microclimate formation and mechanisms of the organism heat exchange.

3. Physiological basics of human heat exchange and thermoregulation, physiological reactions in the comfortable or uncomfortable (hot or cold) microclimate. Influence of high and low temperature with high humidity on the human organism.

4. Acclimatization and adaptation as a complex socio-biological process of human adaptation to the new environment. Physiological changes in the human body, developing during the acclimatization and adaptation to unusual conditions.

5. Definition of «microclimate» and factors, which influence its formation.

6. Research methods and hygienic assessment of microclimate parameters in residential premises and medical institutions (temperature, humidity).

Notes and problems solving

								1																			
																											1
<u> </u>																											<u> </u>
																						 					<u> </u>
<u> </u>																											
																											<u> </u>
<u> </u>																											
																						 					<u> </u>
-		-	-							-		-				-					 -		-	-	-	 \vdash	
<u> </u>																										 \vdash	_
]	
<u> </u>																										\vdash	├
																						 					<u> </u>
<u> </u>																											
	-				-	-			-												 	 					-
																											<u> </u>
<u> </u>																											—
																						 					<u> </u>
<u> </u>					-	-			-												 					 \vdash	
<u> </u>								<u> </u>																		 \vdash	_
<u> </u>					-	-			-												 					 \vdash	
L	<u> </u>		<u> </u>				<u> </u>					 <u> </u>		<u> </u>	<u> </u>	<u> </u>	 \square	<u> </u>									
																										7	_
<u> </u>																											
<u> </u>																										\vdash	
L	<u> </u>				<u> </u>	<u> </u>			<u> </u>																		
L		L	L					L								L							L		L		L
<u> </u>	-	-	-		-					-		-	-	-		-		-	-		 -		-	-	-	 \vdash	
<u> </u>		<u> </u>	<u> </u>					<u> </u>		<u> </u>		<u> </u>				<u> </u>					 <u> </u>		<u> </u>	<u> </u>	<u> </u>	 \vdash	_
																											1
																											<u> </u>
<u> </u>																										\vdash	├

Student's signature_____

Teacher's signature_____

PHYSICAL PROPERTIES OF ATMOSPHERIC AIR. RESEARCH METHODS AND HYGIENIC ASSESSMENT OF MICROCLIMATE PARAMETERS IN RESIDENTIAL PREMISES AND MEDICAL INSTITUTIONS (AIR VELOCITY, ATMOSPHERIC PRESSURE)

Date

Learning objectives:

1. To master the hygienic evaluation of windrose.

2. To learn value of active prevention, taking into account the influence of weather conditions on the human body.

3. To learn physiological reactions to changes in atmospheric pressure

4. To master methods of determination and hygienic assessment of air velocity and atmospheric pressure.

THEORETICAL SECTION

Give a definition

A wind rose — _____

Fill the table

Factor	Health effects
High atmospheric pressure	
Low atmospheric pressure	

PRACTICAL WORK

1. Hygienic assessment of air velocity

The air velocity is measured by _______Standard (normal) values = _______Actual values = _______Conclusion: ______

2. Hygienic assessment of atmospheric pressure

PROBLEMS SOLVING

– Southwest wind — 5 %;

– Western wind — 6 %;

Task. The frequency of wind direction in the region in 5 years is:

- Northwest wind 19 %;
 North wind 9 %;
 Southeast wind 31 %;
 Southern wind 6 %;
- Northeast wind 8 %;
- -East wind 11 %;
- $-\operatorname{Calm} 5\%$.

Draw a wind rose. Indicate optimal location for chemical factory.



TRAINING TEST

1. Choose the parameteres of microclimate:

a. temperature;	c. air movement speed;
b. humidity;	d. light.

2. Choose the correct formula: Deficiency of saturation:

3. Optimal value for humidity in the living	ng room is:
a. 30–65 %;	c. 20–50 mm Hg; e. 40–60 %.
b. 20–50 %;	d. 45–55 %;
4. Air velocity is measured by	
a. Anemometer;	c. Psychrometer;
b. Kata thermometer;	d. Thermometer.
5. Humidity is measured by	
a. Anemometer;	c. Psychrometer;
b. Katathermometer;	d. Thermometer.
6. Temperature is measured by	
a. Anemometer;	c. Psychrometer;
b. Katathermometer;	d. Thermometer.
7. Types of humidity are:	
a. relative;	c. maximal;
b. absolute;	d. barometric.
8. Decompression disease caused by	
a. rapid decrease in the pressure;	c. rapid decrease in the tempetature;
b. rapid increase in the pressure;	d. water pollution.
9. Optimal value for humidity in the livit	ng room is:
a. 30–65 %;	c. > 0.3 m/s; e. 40–60 %.
b. 0.5–1 m/s;	d. < 0.3 m/s;

KEY QUESTIONS

1. The notion of seasonal and meteodiseases. Meteodependence. The value of active prevention, taking into account the influence of weather conditions on the human body.

2. Physiological and hygienic significance of atmospheric pressure. Physiological reactions to changes in atmospheric pressure.

3. Low and high atmospheric pressure, its influence on the human organism. Decompression and altitude diseases

4. Physiological and hygienic significance of air movement. Windrose, methods of it's preparation and hygienic evaluation.

5. Research methods and hygienic assessment of microclimate parameters in residential premises and medical institutions (air velocity, atmospheric pressure).

						1 11	Jie	5 ai	IG]	pro	1113	30	1 1 11	16						

Student's signature_____

Teacher's signature_____

Notes and problems solving

PHYSICAL PROPERTIES OF ATMOSPHERIC AIR. HYGIENIC ASSESSMENT OF THE COMBINED ACTION OF METEOROLOGICAL FACTORS ON THE HUMAN BODY

Date

Learning objectives:

1. To master hygienic evaluation of complex action of meteorological factors on the human body.

2. To learn Kata thermometry and Effective temperature methods, methods of research and hygienic assessment of the thermal state of the human body.

3. To learn measurement indicators characterizing the body's response to the impact of meteorological factors: heatfeelings, average body temperature, cold test, potassium iodide by the method of Minor.

THEORETICAL SECTION

Fill the table

Thermal balance of the human body

Heat gain	Heat loss

Give a definition

Effective temperature (ET) — _____

Thermal comfort — _____

PRACTICAL WORK

Fill in the table «Protocol of hygienic evaluation of complex action of meteorological factors» results of the calculations (p. 12).

Definition of air movement speed with the help of a Kata thermometer

A kata thermometer is heated up to a certain temperature (above air temperature), and then in case of cooling under the influence of temperature and air movement the device will lose certain quantity of heat at a certain level of temperature. Knowing the amount of cooling and the air temperature, we may calculate the air movement speed according to the empirical formulas and tables.

A spherical kata thermometer is placed in a vessel with hot water at the temperature of 65-70 °C until the painted spirit fills half of the top tank. After that the kata thermometer is wiped

dry rag and hanged up on a support in a place where it is necessary to define air movement speed. Further by means of a stop watch time we define the time in seconds for which the column has fallen from T_1 to T_2 . It is possible to take intervals from 40–33 °C, from 39–34 °C, from 38–35 °C, i. e. such an interval, a partial value of which from division of the sum T_1 and T_2 should be equal to 36.5 °C. The test is repeated 2–3 times and the average indexes are calculated. Further on their basis we calculate the amount of cooling H. The cylindrical kata thermometer amount of cooling and the spherical kata thermometer are calculated with an interval of 38–35 °C applying following formulas:

1. H (cooling capacity of the air) = F/a

F — the katathermometer factor, a constant showing the quantity of heat, lost from 1 cm² of the device surface during its cooling from 38 to 35 °C (indicated on the back surface of every katathermometer), mcal/cm²; a — time of cooling, s

2. V (air movement speed) = $\left[\frac{H/Q-0.2}{0.4}\right]^2$

H — cooling capacity of the air; \tilde{Q} — difference between an average body temperature 36.5° and surrounding air temperature

$$Q = T^{\circ}_{body} - T^{\circ}_{air}$$

3. Define effective temperature on the scale and the nomogramm and to give recommendations about its optimization.

Assessment of a human body reactions on the meteorological factors action

1. Skin temperature:

To estimate its dynamics, it should checked strictly in certain points. Which are as follows:

- on the forehead — between superciliary arches on 0.5 cm above their upper edge;

- on the breast — at a breast upper edge;

- on the hand — back surface, between the bases of the first phalanxes of the thumb and index fingers.

If we feel comfort the temperature of a skin of a forehead and a breast is equal to 31-34 °C, a hand skin — 30-31 °C.

2. Definition of the average temperature:

 $T = 0.07 \times F + 0.5 \times B + 0.43 \times H$

F — temperature of the forehead; B — temperature of the breast; H — temperature of the hand; Comfort condition — 32-34 °C

3. Definition of breast and hand surface temperature fluctuations range.

In comfortable conditions the difference between the breast and hand temperature should be 3-4 °C if it is less than 2,5 °C — is a thermal discomfort, and if it is more than 4 °C — it is cold discomfort.

4. Cold Test:

For supervision we choose the limited sites of a skin on an open part of a body and on the usually covered clothes parts (a back, a leg). On the chosen site of skin you take the electrothermometer skin temperature. A metal jar filled with ice is placed on this place for 20–30s. After the expiry term the jar is removed. Each 2 min it is necessary to take skin temperature until the temperature of a skin reach the initial value.

In case the skin temperature will return until the initial level within 5 minutes, it testifies of good adaptation to cold, 10 min — satisfactory, 15 min and more — unsatisfactory.

5. Investigate the function of sweating (the iodinestarch test)

A small area of the skin (forehead) is covered with a solution containing iodine (10 g of castor oil, 15 g of 10 % iodine tincture and 75 ml of ethyl alcohol). After drying, the oiled place is powdered with potato starch. Sweat droplets in the presence of starch with iodine give a blue coloration.

The comfort zone corresponds to the conditions under which perspiration is expressed in the form of individual small dots. The appearance of blue spots indicates uncomfortable conditions for the accumulation of heat in the body.

Parameter	Actual value	Comfort value
Average temperature indoors (°C)		
Relative humidity (%)		
Air speed movement (m/sec)		
Cooling capacity (mkal/cm ² × sec)		
Effective temperature (°ET)		
Physiological reactions of the orga	nism	
Heat sensation character		comfort
Pulse rate		60–80
Body temperature		36 °C
Forehead skin temperature		31–34 °C
Breast skin temperature		31–34 °C
Hand temperature skin		30–31 °C
The average skin temperature		32–34 °C
Range of breast and hand surface temperature fluctuation		3–4 °C
Cold test		Up to15 min
Character of sweating		
Conclusion:		

Protocol of hygienic evaluation of complex action of meteorological factors

PROBLEMS SOLVING

Example. The speed of the air movement in the classroom was determined using a kata thermometer. The factor of the cylindrical kata thermometer equal 588 Mcal/cm², cooling time is 112 sec. In the classroom the temperature is +19 °C.

Find the cooling capacity of the air (H), air movement speed.

Compare the result with the hygienic standard.

1. $H = F/a = 588/112 = 5.25 \text{ mkal/cm}^2 \times \text{sec.}$

2.
$$Q = T^{o}_{body} - T^{o}_{air} = 36.5 - 19 = 17.5^{o}$$
.

3. V =
$$\left[\frac{H/Q-0.2}{0.4}\right]^2 = \left[\frac{5.25/17.5-0.2}{0.4}\right]^2 = 0.0625$$
 m/s.

Conclusion: During the hygienic assessment of the microclimate parameters, it was found that the actual cooling capacity of the air less the norm (actual = $5.25 \text{ mkal/cm}^2 \times \text{sec}$, normal = $5.5-7.0 \text{ mkal/cm}^2 \times \text{sec}$), hot discomfort, air movement speed less the norm (actual = 0.0625 m/s, normal = 0.1-0.2 m/s).

Tasks. 1. In the living room the temperature of dry thermometer is +19 °C, the temperature of wet thermometer is +15 °C (measured by means of an August stationary psychrometer). The factor of the cylindrical kata thermometer equal 533 Mcal/cm², cooling time is 65 seconds.

Calculate the cooling capacity of the air (H), air movement speed and effective temperature (ET). Perform hygienic assessment of calculated data (compare the result with the hygienic standard). If the effective temperature is not included in the line comfort, show how to modify the parameters of the microclimate that ET went into it.

2. Perform a hygienic assessment of the air state in the classroom of the department before the start of classes and at the end of the working day according to the data given in the table:

Investigated indicators	Before the start of classes	At the end of the working day
Air temperature, °C	21.5	26.4
Relative humidity, %	57	78
Cooling capacity of the air	6.5	3.6
Air movement speed, m/s	0.34	0.15

TRAINING TEST

1. The heat from the body can be lost through:

a. conduction; c. convection;

b. evaporation; d. radiation.

2. Methods of assessment complex effect of meteorological factors on the human body:

a. assessment of actual nutrition;

- b. kata thermometry;
- c. method of effective temperatures;
- d. assessment the level of physical development.

3. The effective temperature combines the effects of the following:

a.	air temperature;	c. air velocity;
1.	- 41	1

b. athmospheric pressure; d. relative humidity.

4. Standard (normal value) for effective temperature is:

a. 5.5–7.0 mcal/cm ² × sec;	c. 20–22 °C;
b. 17.2–21.7 °ET;	d. 0.1–0.25 m/sec.

5. Parameters considered in integrated assessment of microclimate by kata thermometry:

a. pressure;	c. air velocity;	e. temperature.
b. sunlight;	d. humidity;	

6. Standard (normal value) for cooling capacity of the air is:

a. 5.5–7.0 mcal/cm ² × sec;	c. 20–22 °C;
b. 17.2–21.7 °ET;	d. 0.1–0.25 m/sec.

7. Parameters that are not taken into account in integrated assessment of microclimate by method of effective temperatures:

a. atmospheric pressure;	c. infrared heat;
b. air speed;	d. temperature.

KEY QUESTIONS

1. Hygienic evaluation of the complex influence of meteorological factors on the human body: methods and their comparative characteristics.

2. What is the kata thermometry method?

3. What is the method of Effective temperatures?

4. Measurement indicators characterizing the body's response to the impact of meteorological factors.

5. Hygienic evaluation of the complex influence of meteorological factors on the human body: methods and their comparative characteristics.

Notes and problems solving

-	1	-	-		1	1	1	1	1		_	 -		-	_	 -	-	-			 -	-	-	-	-	-	-	<u> </u>		1
<u> </u>																														
<u> </u>																													\vdash	
						<u> </u>			<u> </u>												 								\vdash	-
																					 									
																					 								\vdash	
																														<u> </u>
																														_
<u> </u>												 									 								\vdash	
<u> </u>												 									 								\vdash	
																					 									<u> </u>
																														<u> </u>
-			-					-				-				-						-		-	-	-	-			
\vdash					-	-	-	-	-																				\vdash	
<u> </u>					-		-	-				 				 					 								—-	
<u> </u>					-		-	-																					\vdash	├──
																														<u> </u>
								<u> </u>																						
\vdash								-				-																		<u> </u>
⊢	\vdash		-	-	-	-	-	\vdash	-	-		-	-			-			-	-		-		-	-	-	-		\vdash	⊢
					-		-																						<u>├</u>	

Student's signature_____

Teacher's signature_____

PHYSICAL PROPERTIES AND CHEMICAL COMPOSITION OF ATMOSPHERIC AIR. FINAL CLASS

CONTROL QUESTIONS

1. Temperature of air characteristics, influence on a human organism, research methods, hygienic assessment.

2. Humidity of air characteristics, influence on a human organism, research methods, hygienic assessment.

3. Barometric pressure of air characteristics, influence on a human organism, research methods, hygienic assessment.

4. Movement of air characteristics, influence on a human organism, research methods, hygienic assessment. Wind rose.

5. Hygienic assessment of complex influence of meteorological factors on the human body: methods and their characteristics.

6. Heat exchange with the environment. Consequences of its violations: overheating, overcooling. Preventive measures.

7. Climate and weather. Acclimatization. Hygiene measures to facilitate acclimatization. Meteotropic diseases and their prevention.

EXAMPLES OF CONTROL CASE TASKS

1. Indications of dry and wet bulbs of aspirating August psychrometer in the living room are respectively +19 °C and +15 °C. Atmospheric pressure is 755 mmHg.

Calculate absolute humidity of air, relative humidity of air, saturation deficiency and dewpoint temperature. Perform hygienic assessment of calculated data (compare the result with the hygienic standard).

2. In the classroom the temperature of dry thermometer is +18 °C, the temperature of wet thermometer is +15 °C (measured by means of an August stationary psychrometer). The factor of the cylindrical katathermometer equal 535 mkal/cm², cooling time is 65 sec.

Calculate the cooling capacity of the air (H), air movement speed and effective temperature (ET). Perform hygienic assessment of calculated data (compare the result with the hygienic standard). If the effective temperature is not included in the line comfort, show how to modify the parameters of the microclimate that ET went into it.

Northwest wind	11 %
North	6 %
Northeast wind	5 %
East wind	8 %
Southeast wind	30 %
Southern wind	15 %
Southwest wind	7 %
The western wind	15 %
Calm	3 %

3. The frequency of wind direction in the region for 5 years.

Draw a wind rose. Indicate optimal location for oil refineries.

HYGIENIC ASSESSMENT OF THE IMPACT OF PLACEMENT CONDITIONS ON HUMAN HEALTH. RESEARCH METHODS AND HYGIENIC ASSESSMENT OF NATURAL LIGHTING IN ENCLOSED PREMISES

Date

Learning objectives:

1. To learn the hygienic requirements for natural lighting in different premises.

2. To master the geometrical, lighting engineering methods of natural lighting indices determination, to learn how to assess the results of instrumental measuring, and to draw a hygienic conclusion about natural lighting in differing premises.

THEORETICAL SECTION

Fill the table

Effects of separate components of solar radiation on organism. Spectral distribution and biological activity of solar radiation

Kind of radiation	Wave length	Biological affect
Infrared		
Visual		
Ultraviolet:		
Range A		
Range B		
Range C		

Give a definition

Coefficient of natural light — _____

Light coefficient — _____



PRACTICAL WORK

Hygienic estimation of natural lighting

The geometrical method

1. Determination of the light coefficient (LC)

LC (light coefficient) = $S_w \times n : S_f$

 S_w — glazed area of 1 window; n — number of the windows; $S_{\rm f}$ — floor area

Standard (normal) values = _____

Actual value = _____

Conclusion:

2. Determination of the depth coefficient (DC)

DC (depth coefficient) = D/h D — room depth; h — the height of the upper edge of the window from the floor Standard (normal) values = _____

Actual value = _____ Conclusion: _____

3. Determination of the light and aperture angles (α,β)

Standard (normal) values = _____





Fig. 1. Light angle: AB — is the distance from the working place to the lower line of the window; BC — is the height of the window; BC/AB — tg of angle α . Using the table, transform the tg into the angle value Fig. 2. Aperture angle:

BD — distance between the line on the window glass from the opposite subject and the lower line of window; BD/AB — is tg of angle γ ; Using the table, transform the tg into the angle value angle of opening β = angle α – angle γ

Actual value = _____

Conclusion:

The technical lighting method

Device for measurement _____

Determination of the coefficient of natural light (CNL)

CNL (coefficient of natural light) = $E_1/E_2 \times 100$ %

 E_1 — percentage of light exposure of a point indoors; E_2 — light exposure of the external point taking place (outdoor illuminance)

Standard (normal) values = _____

Actual value = _____

Conclusion:

PROBLEMS SOLVING

Example. On a workplace in a university laboratory illumination was 360 lux. External lighting outdoor was 16300 lux. What is the coefficient of natural lighting? Compare the result with the hygienic standard.

CNL (coefficient of natural light) = $E_1/E_2 \times 100 \% = 360/16300 \times 100 \% = 2.2 \%$ Conclusion: During the hygienic assessment of the natural lighting, it was found that the coefficient of natural light less the norm (actual = 2.2 %, normal = $\ge 2.5 \%$). **Example.** Depth of the living room -7 m, length -7 m, height -3 m. The room has two windows, their height is 2.3 m above the floor, a glazed area of each of them 2.7 m². Calculate the coefficient of light, coefficient of depth. Compare the result with the hygienic standard.

 $LC = Sw : Sf = 2.7 \times 2 : 7 \times 7 = 5.4 : 49 = 1 : 9$

DC = Depth of the room/height of windows above the floor = 7/2.3 = 3

Conclusion: During the hygienic assessment of the natural lighting, it was found that the light coefficient is optimal (actual = 1 : 9, optimal = 1 : 8 - 1 : 10), coefficient of depth not normal (actual = 3, optimal = ≤ 2.5).

Task. Depth of the living room — 5.5 m, length — 3.5 m, height — 3.2 m. The room has two windows, their height is 2.8 m above the floor, a glazed area of each of them 2.6 m². Internal natural lighting is 360 lux, lighting outdoor — 15300 lux.

Calculate the coefficient of natural light, the light coefficient, coefficient of depth. Compare the result with the hygienic standard.

KEY QUESTIONS

1. Significance of solar radiation, the spectral structure of solar light.

2. Concept of solar starvation, reasons, manifestations and prevention.

3. Physiological effect of various components of a solar spectrum.

4. Hygienic requirements to natural lighting, factors which influence the natural lighting.

5. The natural lighting determination methods.

6. Light coefficient and the method of its determination.

7. The light angle, the aperture angle: hygienic standards and the methods of determination.

8. The coefficient of natural illumination, method of its determination, hygienic standards for different rooms.



Notes and problems solving

Student's signature_____

Teacher's signature

HYGIENIC ASSESSMENT OF THE IMPACT OF PLACEMENT CONDITIONS ON HUMAN HEALTH. RESEARCH METHODS AND HYGIENIC ASSESSMENT OF ARTIFICIAL LIGHTING IN ENCLOSED PREMISES

Date

Learning objectives:

1. To learn the role and the meaning of the rational artificial illumination as the means of lengthening the activity period of people, and disease and fatigue prevention.

2. To master the methods of the measurement and hygienic assessment of artificial illumination in different premises with the help of a luxmeter and calculation methods.

THEORETICAL SECTION

Requirements for artificial lighting:



Fill the table

Characteristics of artificial lighting sources

	Incandescent lamps	Fluorescent lamps
Advantages		
Disadvantages		

PRACTICAL WORK

Hygienic assessment of artificial illumination by the «Watts» method

 $E = P \times Et / 10 \times K$,

K = 1,3 (reserve coefficient for residential and public premises); $P = power of one lamp \times number of lamps in the room / S_f; Et = for luminescent lamps = 150 lux; for incandescence lamps — in table «Size of the minimum horizontal light (lux) at specific capacity 10 W/m²»$

Actual values = _____

Conclusion: _____

PROBLEMS SOLVING

Example. Room in a hostel area of 24.4 m^2 illuminated by 3 incandescent lamps of 150 watts each, light — direct. Calculate the value of illumination in lux. Compare the result with the hygienic standard.

Et = 46 lux (from the table «Size of the minimum horizontal light (lux) at specific capacity 10 W/m²»)

 $P = power \ of \ one \ lamp \times number \ of \ lamps / \ S_f = 150 \times 3$ / 24.4 = 18.44 W/m^2

 $E = P \times Et / 10 \times K = 18.44 \times 46 / 10 \times 1.3 = 848.24 / 13 = 65 lux$

Conclusion: During the hygienic assessment of the artificial illumination, it was found that the artificial illumination is sufficient (actual = 65 lux, optimal = 50 lux and more).

Example. Reading hall area of 150 m^2 illuminated by 35 luminescent lamps of 45 watts each. Calculate the value of illumination in lux. Compare the result with the hygienic standard.

Et = 150 lux (constant)

 $P = power \ of \ one \ lamp \times number \ of \ lamps / S_f = 45 \times 35 \ / \ 150 = 10.5 \ W/m^2$

 $E = P \times Et / 10 \times K = 10.5 \times 150 / 10 \times 1.3 = 121.2 lux$

Conclusion: During the hygienic assessment of the artificial illumination, it was found that the artificial illumination is insufficient (actual = 121.2 lux, optimal = 400 lux and more).

Tasks. 1. Area of room in a hostel is 13 m^2 , illuminated by 2 incandescent lamps of 75 watts each, light — reflected. Calculate the value of artificial light using the method of «Watt», compare it with the hygienic standards for the living room.

2. Reading hall area of 150 m^2 illuminated by 26 luminescent lamps of 40 watts each. Calculate the value of illumination in lux, compare it with the hygienic standard.

TRAINING TEST

1. Coefficient of natural light

a. $CNL = E_1 / E_2 \times 100 \%;$	c. $LC = S_w:S_f;$	e. tg $\beta = h_2 / L_1 + L_2$.
b. $E = 150 P/13;$	d. $E = P \times Et / 10 \times K;$	

2. The light coefficient is:

a. the ratio of illumination in the open air to illumination indoors;

b. the ratio of light angle to aperture angle;

c. the ratio of the area of windows to the floor area;

d. the ratio of window area to tangent of aperture angle.

3. Compared to incandescent lamps, fluorescent lamps have several advantages:

a. are characterized by low brightness;

b. create diffused light that does not give harsh shadows;

c. do not have a glare;

d. the appearance of monotonous noise during operation.

4. Identify the range of ultraviolet radiation which promotes formation of vitamin D (in nanometers):

a. 265–315;	c. 200–280;	e. 280–320.
b. 280–310;	d. 315–400;	

5. Types of lighting:

a. domestic; b. climatic; c. natural; d. artificial.

KEY QUESTIONS

1. Hygienic requirements to artificial lighting.

2. Sources of artificial lighting and their hygienic characteristics.

3. Systems of artificial lighting.

4. Methods of measurement of artificial lighting.

Notes and problems solving

Student's signature_____

Teacher's signature_____

HYGIENIC ASSESSMENT OF THE IMPACT OF PLACEMENT CONDITIONS ON HUMAN HEALTH. NATURAL AND ARTIFICIAL VENTILATION, TYPES AND THEIR HYGIENIC CHARACTERISTICS. RESEARCH METHODS AND HYGIENIC ASSESSMENT OF CARBON DIOXIDE CONCENTRATION. HYGIENIC ASSESSMENT OF VENTILATION INDICATORS: AIR CUBE, VENTILATION VOLUME, AIR EXCHANGE RATE

Date_____

Learning objectives:

1. To learn the role and hygienic value of the ventilation.

- 2. Studying types of ventilation their characteristics.
- 3. Studying indicators of the effectiveness of ventilation of premises.
- 4. To learn the research methods and hygienic assessment of carbon dioxide concentration.

5. To learn the hygienic assessment of ventilation indicators.

THEORETICAL SECTION

Give a definition

Ventilation — _____

Effects of Air Pollution:

1	
2.	
3	
4.	
5.	
6.	
7.	

Role of air in human health:

1	 	 	
2.			
3.			
4.			
5.			

Fill the table

Sources of atmospheric air pollution

Outdoor Air Pollution	Indoor Air Pollution

Give a definition

Maximu	m permissible concentration (MPC) —
Ventilat	on volume —
Frequen	cy rate of air exchange in 1 hour —
Types of	ventilation:
1	
a.	
b	
c.	
2 a. b. c.	
Advanta 1	ges of mechanical ventilation are:
2	
3	
4	
5	
J	

PRACTICAL WORK

Calculating necessary ventilation volume and necessary frequency rate of air exchange in 1 hour

1. L_n (necessary ventilation volume) = $\frac{C \times n}{P_1 - P_2}$

C — volume of CO₂, expired by one person per hour, 22.6 l/h; n — number of people indoors; P₁ — CO₂ maximum admissible concentration in pro mil (1.0‰ = 0.1 %); P₂ — CO₂ concentration in the atmospheric air in pro mil (0.4‰ = 0.04 %)

Actual value = _____

2. K_n (necessary frequency rate of air exchange in 1 hour) = L_n / V V — indoor cubage (cubature), m³

Actual value = _____

PROBLEMS SOLVING

Problem solving algorithm:

1. L_n (necessary ventilation volume) = $\frac{C \times n}{P_1 - P_2}$

2. K_n (necessary frequency rate of air exchange in 1 hour) = L_n / V

3. L_a (actual ventilation volume) =

a) L_a =
$$\frac{C \times n}{P - P_2}$$

 $P - CO_2$ actual concentration in a premise in pro mil

b) $L_a = A \times B \times C = S \times v \times t$

A (S) — ventilation apertures area (window leaf), m^2 ; B (v) — speed of the air movement through ventilating aperture, m/sec; C (t) — time of aeration, sec

4. K_a (actual frequency rate of air exchange in 1 hour) = L_a/V

5. Compare necessary and actual values, make a conclusion

Example. During the sanitary and hygienic examination of the air in the postoperative ward of the surgical department (indoor cubage 69.7 m³), where 4 patients are located, the CO₂ content was 0.12 %. Determine the necessary and actual volume of ventilation and the rate of air exchange.

Determine the required ventilation volume and air exchange rate:

1. L necessary $(L_n) = (C \times n) / (P_1 - P_2) = (22.6 \times 4) / (1 - 0.4) = 150.67 \text{ m}^3$

2. K necessary $(K_n) = L_n / V = 150.6 / 69.7 = 2.16$

Determine the actual ventilation volume and the rate of air exchange:

3. L actual (L_a) = (C × n) / (P – P₂) = (22.6 × 4) / (1.2 – 0.4) = 113 m³

4. K actual = $L_a / V = 113 / 69.7 = 1.62$

Conclusion: the actual ventilation volume (113 m^3) is significantly lower than the required ventilation volume (150.67 m^3) , the actual frequency of air exchange (1.62 times) is significantly lower than the required one (2.16 times), which indicates the ineffectiveness of ventilation in the examined ward. It is necessary to increase the ventilation time.

Example. In a room with an indoor cubage of 60 m³, where there are 3 people, ventilation takes place due to a window, which is opened for 10 minutes every hour. The air velocity in the ventilation opening is 1 m/s, the window area is 0.15 m^2 . Calculate the ventilation volume, frequency rate of air exchange in 1 hour. Compare the result with the hygienic standard.

Determine the required ventilation volume and air exchange rate:

1. L necessary $(L_n) = (C \times n) / (P_1 - P_2) = (22.6 \times 3) / (1 - 0.4) = 113 \text{ m}^3$

2. K necessary $(K_n) = L_n / V = 113 / 60 = 1.88$

Determine the actual ventilation volume and the rate of air exchange:

3. L actual (L_a) = S × v × t = 0,15 × 1 × 600 = 90 m³

4. K actual = $L_a/V = 90/60 = 1.5$

Conclusion: the actual ventilation volume (90 m^3) is significantly lower than the required ventilation volume (113 m³), the actual frequency of air exchange (1.5 times) is significantly lower than the required one (1.88 times), which indicates the ineffectiveness of ventilation in the examined ward. It is necessary to increase the ventilation time.

Tasks. 1. Calculate the ventilation volume, frequency rate of air exchange in 1 hour for the living room of $4.5 \times 5.5 \times 3.0$ m, which is home to 4 people. The concentration of CO₂ in the moment of research was 0.13 %. Compare the result with the hygienic standard. Make a hygienic assessment of the effectiveness of ventilation and air exchange rate.

2. There are 3 people in the room of $3.5 \times 2.5 \times 3$ m. Ventilation of the room is realized by opening the window for 10 minutes every hour. Area of the window opening — 0.5×0.3 m², air speed — 1.05 m/s. Calculate the ventilation volume, frequency rate of air exchange in 1 hour. Compare the result with the hygienic standard. Make a hygienic assessment of the effectiveness of ventilation and air exchange rate.

TRAINING TEST

1. Indoor pollutants include:

a. ozone;	c. radon;
b. tobacco smoke;	d. chlorides;
e. chemicals, released from household	cleaners.

2. Air quality in living rooms and public buildings assessed by content:

a. ozon	e;	c. carbon dioxide;	e. oxygen.
b. argo	n;	d. nitrogen;	
3. Natural ver	ntilation dep	ends on:	
a. effec	ts of differen	d. diffusion of metals;	
b. diffu	sion of gases	s;	e. water diffusion.
c. perfl	ation and asp	piration;	
4. Systems of	ventilation:		
a. artifi	cial;	c. chemical;	e. permanent.
b. accid	dental;	d. natural;	

KEY QUESTIONS

- 1. Indoor air environment.
- 2. Sources of air pollution of residential and public premises.
- 3. Indicators of indoor air cleanliness.
- 4. Ventilation, its hygienic value.
- 5. Types of ventilation and their characteristics.
- 6. Indicators of the effectiveness of ventilation of premises.

7. Sources of air pollution and hygienic requirements for ventilation in the premises of healthcare organizations. Criteria for the cleanliness of the air of the premises of healthcare organizations.

8. The role of ventilation in the prevention of nosocomial infection.9. Hygienic requirements for the organization of air exchange in the ward. Ventilation and aeration mode.

				1		-	1					1									1				<u> </u>	
	 									 						 							 			 <u> </u>
																										1
<u> </u>	 			<u> </u>			<u> </u>			 		<u> </u>				 							 		\rightarrow	
																										1
<u> </u>	 									 						 							 		\rightarrow	 <u> </u>
<u> </u>										 															-+	 <u> </u>
	 									 																 <u> </u>
<u> </u>										 															-+	 <u> </u>
																										1
		L				L																				 \vdash
																										<u> </u>
<u> </u>	 									 						 							 			 <u> </u>
<u> </u>	 									 						 							 		\rightarrow	 <u> </u>
										 							_		_							
	 									 						 							 			
										 							_		_							
										 																
																									-+	
		L	L	L		L	L				L	L														
		<u> </u>	<u> </u>			<u> </u>			\square		<u> </u>							\mid		 \mid					$ \rightarrow $	
			-					\vdash			-			-	-	-	-		-	 		-	-			
		L				L																				 \vdash
																										<u> </u>
																									\neg	
$\left - \right $																		$\left - \right $								 ┢──┤
																									-+	
			<u> </u>								<u> </u>														$ \rightarrow$	
																	_		_						-+	
																										l l
	 				-					 			-			 		 ·				_	 			

Notes and problems solving

Student's signature_____

Teacher's signature_____

HYGIENIC ASSESSMENT OF THE IMPACT OF PLACEMENT ON HUMAN HEALTH. FINAL CLASS

CONTROL QUESTIONS

1. Natural lighting of dwellings: hygienic significance, criteria and evaluation methods.

2. Significance of solar radiation, the spectral structure of solar light. Physiological effect of various components of a solar spectrum: UV, visible beams, IR-rays

3. Concept of solar starvation, reasons, manifestations and prevention.

4. Hygienic requirements to natural lighting. Factors which influence the natural lighting. Determination methods

5. Prevention of the vision disorders.

6. Artificial lighting: hygienic significance, research methods.

7. Comparative hygienic assessment of various sources of artificial lighting.

8. Indoor air environment.

9. Sources of air pollution of residential and public premises.

10. Indicators of indoor air cleanliness.

11. Ventilation, its hygienic value.

12. Types of ventilation and their characteristics. Indicators of the effectiveness of ventilation of premises.

13. Sources of air pollution and hygienic requirements for ventilation in the premises of healthcare organizations. Criteria for the cleanliness of the air of the premises of healthcare organizations.

14. The role of ventilation in the prevention of nosocomial infection.

15. Hygienic requirements for the organization of air exchange in the ward. Ventilation and aeration mode.

EXAMPLES OF CONTROL CASE TASKS

1. On a workplace in a university laboratory illumination was 320 lux. External lighting outdoor was 15200 lux. What is the coefficient of natural lighting? Compare the result with the hygienic standard.

2. Depth of the room — 6 m, length — 5 m, height — 3 m. The room has two windows, their height is 2.5 m above the floor, a glazed area of each of them 2.5 m². Calculate the coefficient of light, coefficient of depth. Compare the result with the hygienic standard.

3. Center of workplace is 3 m from the window. The height of the upper edge of the glazing windows from the horizontal plane workplace 1.5 m. 7.5 m from the window there is a nearby building that rises above 5 m from the horizontal plane. Calculate the light angle on the workplace and the angle of the aperture (with picture). Compare the result with the hygienic standard.

4. Room in a hostel area of 10.3 m^2 illuminated by 2 incandescent lamps of 100 watts each, light — direct. Calculate the value of illumination in lux. Compare the result with the hygienic standard.

5. Reading hall area of 195 m² illuminated by 40 luminescent lamps of 50 watt each. Calculate the value of illumination in lux. Compare the result with the hygienic standard.

6. Calculate the actual and necessary number of air for the living room of $4.2 \times 4.1 \times 2.7$ m, which is home to 4 people. The concentration of CO₂ in the moment of research was 0,18 %. Calculate the ventilation volume, frequency rate of air exchange in 1 hour. Compare the result with the hygienic standard.

7. In the lab with $4.9 \times 5.7 \times 3.2$ m work 5 people. Dimensions of ventilation openings 0.15×0.4 m, air speed 1.3 m/s. Duration of ventilation is 10 minutes. Calculate the ventilation volume, frequency rate of air exchange in 1 hour. Compare the result with the hygienic standard.

HYGIENIC ASSESSMENT OF DRINKING WATER QUALITY. RESEARCH METHODS AND HYGIENIC ASSESSMENT OF PHYSICAL, ORGANOLEPTIC AND CHEMICAL INDICATORS OF WATER QUALITY

Date_____

Learning objectives:

1. To learn methods of research and hygienic assessment of the physical and organoleptic characteristics.

2. To learn methods of research and hygienic assessment of the content of substances in water, indifferent and have a positive physiological significance (carbonates, bicarbonates, calcium, magnesium).

3. To learn methods of research and hygienic assessment of chemical indicators of organic pollution of water (ammonia nitrogen, nitrite, oxidation).

4. To learn water treatment methods: clarification (coagulation and filtration), disinfection (physical methods: boiling, water treatment with UV, ultrasound; chemical methods: chlorination, ozonation).

5. To analyze advantages and disadvantages of different methods of water disinfection.

THEORETICAL SECTION

Fill the tables

Compare the sources of drinking water

Criteria	Surface water	Ground water

Communicable diseases associated with water

Group	Disease	Pathogene	Source
Waterborne diseases			
Waterwashed diseases			
Water-based diseases			
Water-related diseases			
Waterdispersed diseases			

Give a definition

Hardness —	 	
Oxidability —	 	
Residual chlorine —		

Fill the table

Compare the water supply systems

Individual system	Central system

PRACTICAL WORK

Determination physical and organoleptic properties of water.

Determination of chromaticity of water

100 ml of filtered test water is poured into the cylinder, then look down into the cylinder on a white background and select a cylinder on a standard scale, the color of the solution in which corresponds to the color of the test water (Table 1).

Table 1

	Bia	iuai u s			laticity	utilit	ion			
Cylinder number	1	2	3	4	5	6	7	8	9	10
Chromaticity degrees	0	5	10	15	20	25	30	40	50	60

Standard scale for chromaticity definition

The result:

Conclusion:

Determination of water transparency

The test water is well shaken and poured into a cylinder. A slide is placed under the cylinder, on which the Snellen font is pasted. Then gradually release the water through the tap, find the maximum height of the column of water through which it is possible to read the text of the font. This height of the remaining column of water, indicated in cm, will express the degree of transparency.

The result:_____

Conclusion:

Chemical analysis of water

Qualitative determination of ammonia salts with approximate quantitative assessment (express method)

Reagents:

1. Nessler reagent.

2. Potassium-sodium tartrate (Ferrotic salt), 50 % solution.

Technique of determination: 10 ml of the test water is poured into a test tube with a diameter of 13–14 mm, 0.2–0.3 ml of potassium-sodium tartrate and 0.2 ml of Nessler reagent are added. After 10–15 minutes, an approximate determination is carried out according to Table 2.

Table 2

TPP officient of unmonite und unmonitem for						
	Ammonia and ammonium					
on the side	above	ions, mg/L				
without color	without color	< 0.05				
without color	very pale yellowish	0.1				
very pale yellowish	pale yellowish	0.3				
pale yellowish	yellowish	0.5				
yellowish	light yellow	1.0				
light yellow	yellow	2.5				
yellow	dark-yellow	5				
dark-yellow	brown, muddy	10				

Approximate content of ammonia and ammonium ions

The result:

Conclusion:_____

Qualitative determination of nitrite nitrogen with approximate quantitative assessment (express method)

Reagents: Griss reagent.

Technique of determination: 10 ml of test water is poured into a test tube with a diameter of 13-14 mm, 1 ml of 10 % Griss reagent is added and heated to 50–60 °C in a hot water. After 10 min, the resulting color is compared with the scale of visual determination (Table 3). For the development of color without heating, samples are left for 15-20 min at room temperature.

Approximate content of nitrites

	Color					
on the side	above	Nitrites, ing/L				
without color	without color	<0,003				
without color	very pale pink	0,003				
very pale pink	pale pink	0,007				
pale pink	very light pink	0,013				
very light pink	light pink	0,05				
light pink	pink	0,1				
pink	intensive pink	0,2				
intensive pink	red	0,5				
intensive pink	dark red	1,0				

The result:_____

Conclusion:_____

Determination of water hardness

Reagents:

1. Ammonia-buffer solution.

2. 0.1n. trilon B solution.

3. Indicator: Eriochrome black T.

Technique of determination: 100 ml of pre-filtered test water is poured into the flask, 5 ml of ammonia-buffer solution and 6 drops of the eriochrome black indicator are added, then slowly titrated with 0.1 n. trilon B solution until the wine-red color turns blue.

The hardness of the analyzed water in milligram equivalents will be equal to the number of milliliters of 0.1 n. solution of trilon B, which went to titration of 100 ml of water.

The result:_____

Conclusion:_____

Determination of temporary water hardness

Reagents:

1. 0,1 n. hydrochloric acid solution.

2. Indicator: methylorange.

Technique of determination: 100 ml of the test water is poured into the flask, 2 drops of methylorange are added and titrated from the burette with 0,1 n. hydrochloric acid solution until the yellow color changes to a weak pink; for better capture of the transition of one color to another, a flask with the same water and 2 drops of methylorange in it is placed next to it. The amount of milliliters of 0.1 n. hydrochloric acid solution used for titration of 100 ml of water corresponds to the hardness of water in milligram equivalents.

The result:_____

Conclusion:_____

Determination of the oxidability of water with Mor salt

Reagents:

1. 0.01 n. solution of potassium permanganate (KMnO₄).

2. 0.01 n. solution of Mor salt.

3. 25 % solution of sulfuric acid (H₂SO₄).

Technique of determination: Pour 100 ml of tested water into the flask, add 5 ml of 25 % H_2SO_4 solution and pour 10 ml of 0.01 n. KMnO₄ solution. Wait for 15 minutes. Then add 10 ml of 0.01 n. solution of Mor salt to the sample. The discolored solution should be titrated with potassium permanganate until a weak pink staining appears, the consumption of 0.01 n. KMnO₄ (A₁) solution is recorded. This consumption of permanganate indicates the amount of titrant, which is necessary for the oxidation of organic substances in 100 ml of water.

For the final calculation, it is necessary to set the correction factor (K) KMnO₄. Pour into the same flask 10 ml 0.01 n of the solution of the salt of Mor and again titrate 0.01 n. with a solution of KMnO₄ until a pale pink color appears, the consumption of 0.01 n is recorded solution KMnO₄ (A₂).

The calculation is made according to the formula:

 $X = ((A_1 + A_2) - 10) \times 0.8 \times K$

The result:_____

Conclusion:

KEY QUESTIONS

- 1. Physiological, hygienic and epidemiological importance of water.
- 2. Diseases related to water.
- 3. Sources of drinking water.
- 4. Sources of water pollution.
- 5. Water supply systems.
- 6. Requirements to drinking water.

Notes and problems solving



Student's signature_____

Teacher's signature

HYGIENIC ASSESSMENT OF DRINKING WATER QUALITY. METHODS FOR IMPROVING WATER QUALITY

Learning objectives:

Date_____

1. To learn water treatment methods: clarification (coagulation and filtration), disinfection (physical methods: boiling, water treatment with UV, ultrasound; chemical methods: chlorination, ozonation).

2. To analyze advantages and disadvantages of different methods of water disinfection.

THEORETICAL SECTION

Fill the table

Classification of water treatment methods

Ba	Special	
Clarification an	d decolouration	
Disinf	fection	
Chemical	Physical	

Give a definition

Sedimentation — _____

Coagulation — _____

Advantages and disadvantages of different methods of water disinfection

Method	Chlorination	Ozonation	UV-disinfection			
Advantages						
1 Iu vultuges						

End of the table

Method	Chlorination	Ozonation	UV-disinfection		
Disadvantages					

PRACTICAL WORK

Determination of the amount of aluminum sulfate required for water coagulation

Reagents:

1. 12 % sodium bisulfate solution.

2. Indicator: 1 % alcohol solution of methylroth.

3. Solution of Al₂ (SO₄)₃ × 18 H₂O.

Technique of determination: When water is coagulated with aluminum sulfate, the temporary water hardness is determined first of all. To do this, 100 ml of water is poured into a glass or flask, 2 drops of 1 % alcohol solution of methylroth are added. The water turns yellowish. It is titrated drop by drop 12 % sodium bisulfate solution before pink staining, the number of drops of sodium bisulfate used for titration indicates the temporary hardness of water in degrees.

Having determined the disposable hardness, they proceed to experimental coagulation of water with a solution of Al_2 (SO₄)₃ × 18 H₂O in three glasses. 200 ml of test water is poured into these glasses.

X drops of coagulant are added to the first glass; X - 4 drops are added to the second glass; X - 8 drops are added to the third glass and thoroughly mixed.

The choice of the number X (drops) depends on the temporary water hardness and is carried out according to table:

Temporary water hardness	4°	5°	6°	7°	8°	9°	10°	11°	12°	
X (the amount of drops)	8	12	16	20	24	28	32	36	40	

For example, the temporary hardness is 8° , then 24 drops of coagulant are added to the first glass, 20 drops to the second glass, and 16 drops to the third. At the time of application of the coagulant, the water is mixed for 1 min. Then they are monitored for 10-15 minutes. Choose a glass in which coagulation goes best (large flakes are formed earlier, quickly settling to the bottom). If coagulation is the same in all glasses, then choose the one where the least amount of coagulant is introduced. Based on the selected glass, the required amount is calculated coagulant per 1 liter of water. Suppose we chose the first glass, where 24 drops of aluminum sulfate solution were added. If 24 drops of coagulant are needed for 0.2 liters of water, then 120 drops or 4.8 ml per 1 liter.

The result:

Conclusion:
Determination of the amount of bleach required for chlorination

Technique of determination: To determine the required dose of chlorine during chlorination, a trial chlorination of water is carried out. Field test conditions chlorination is carried out in three cups in each of which is poured 200 ml of the investigated water, put the glass sticks and using calibrated pipettes (25 drops equal 1 ml) add a 1 % chlorine solution: in the first — 1 drop in the second drop 2, the third 3 drops. The water in the glasses is mixed well and after 30 minutes the presence of residual chlorine in it is determined. To do this, 2 ml of a 5 % solution is added to each glass potassium iodide, 2 ml of hydrochloric acid (1: 5), 1 ml of 1 % starch solution and mix thoroughly. In the presence of residual chlorine, the water turns blue, the more intense the more residual chlorine it contains. The water in the glasses where the blue staining appeared is titrated drop by drop with a 0.7 % solution of sodium thiosulfate before discoloration, stirring it after adding each drop.

To calculate the dose, choose the glass where the discoloration occurred from 2 drops of sodium thiosulfate, because the content of the residual chlorine in this glass is 0.4 mg / 1 (1 drop of 0.7% sodium thiosulfate solution binds 0.04 mg of chlorine, which, when converted to 1 liter, corresponds to $0.04 \times 5 = 0.2 \text{ mg} / 1$). If the discoloration occurred from a single drop, the residual chlorine content is not enough — 0.2 mg/ 1; with discoloration from 3 drops, the residual chlorine content is excessive — 0.6 mg / 1. Depending on the results of the trial chlorination, the amount of bleach required for chlorination of 1 liter of water is calculated.

The result:	 	
Conclusion:		

PROBLEMS SOLVING

Task: In urban settlement water supply based on the use of artesian water. Well depth 185 m water disinfection system recently is not functioning. Analysis of water from the tap:

- chromaticity, degrees 20, a yellowish color;
- transparency cm 23;
- smell, points 4, moldy;
- taste, points 3, astringent;
- ammonia nitrogen, mg/l 1.3;
- nitrite nitrogen, mg/l 0;
- nitrate nitrogen, mg/l 22;
- sulphates, mg/l 36;
- chlorides, mg/l 58;
- iron, mg/l 26;
- total hardness, mEq/L 12;
- oxidability, mg $O_2/l 6$;
- the total bacterial count in 1 ml 300;
- total coliform bacteria per 100 ml 80.

1. Specify normative value for each of parameter. Evaluate each parameter has according to the standard value.

2. Make a written report on the suitability of water for drinking purposes.

3. What methods are used for improving the quality of drinking water?

TRAINING TEST

1. The uses and applications for w a. drinking and food preparat b. personal hygiene activities	ater include the follow tion purposes; s including bathing and pet cleaning	wing: I laundering;
2. The main sources of drinking w and	ater are:	
3. Correlate the options:		
1. surface water	2. grou	nd water
a. more sanitary;b. less exposed to pollution;c. more contaminated;d usually clean;e. has low mineralization.		
4. Nature of water pollution:		
a. accidental;	d. diffuse;	
b. agricultural;	e. chemical;	
c. microbiological;	f. permanent.	
5. Waterborne diseases are:		
a. leishnmaniasis;	c. hepatitis A;	e. trachoma;
b. typhus;	d. salmonellosis;	f. typhoid.
6. Water-washed diseases are:		
a. scabies;	c. typhus;	e. leishnmaniasis;
b. chicken pox;	d. trachoma;	f. hepatitis A.
7. Water-based diseases are:		
a. COVID-19;		
b. hepatitis D;		
c. dracunculiasis.		
8. Water-related diseases are:		
a. malaria;	c. gangrene;	
b. syphilis;	d. trypanosomiasis.	
9. Methemoglobinemia related to		_in drinking water.
10. Hardness of water may be:		
a. temporary;	d. seasonal;	
b. biological;	e. physical;	
c. permanent;	f. natural.	_
11. Guidelines for bacteriological	water quality are bas	ed on:
a. Samonella typni;	c. Escherichia coli;	
b. Ascaris fullibricoides;	d. Hepatitis A virus	
12. Correlate the options:		
1. Temporary nardness	a. snortage in drinking	g water causes the endemic gotter.
2. Fluoride	o, are toxic substance	by boiling water
J. IOUIIIC A. Heavy metals	d have been linked to	vy voining water.
5 Nitrates	e excess may lead to	dental or skeletal fluorosis
Answer:		contail of Skelotal Hubiosis.

13. Name types of water supplie	es and systems:				
1)	; 2)				
and then correlate the or	otions:				
a. public system;	d. easy to contr	rol;			
b. private system;	e, higher risk of infections in a case of accider				
c. mostly not controlled;	f. epidemiologi	ical risk of disease is lower.			
14. Basic methods of quality wa	ter improving:				
a. clarification;	c. fluoridation;	e. chemical;			
b. degassing;	d. bacterial;	f. organic.			
15. Removal of coarse grained s	uspended particles is	•			
16. Removal of Fe by their oxid	ation and transformation	n to insoluble compounds is called			
· · ·		r			
17. For drinking water disinfect	tion are used:				
a. ozone;	c. chlorine substances;	; e. ionizing radiation;			
b. heavy metals;	d. ultrasound;	f. aluminum.			
18. Organoleptic parameters of	water:				
a. smell;	c. colifags;	e. general coliformed bacteria;			
b. ammonia;	d. chlorine;	f. chromaticity.			
19. Chemical parameters of wat	ter:				
a. chromaticity;	d. nitric acids salts;				
b. chlorides;	e. thermotolerant colif	formed bacteria;			
c. taste	f. general microbic nu	mber.			
20. Microbiological parameters	of water:				
a. smell;	c. oxidability;	e. colifags;			
b. salts of ammonia;	d. transparency;	f. general coliformed bacteria			

KEY QUESTIONS

- Classification of methods to improve water quality (water treatment).
 Methods of clarification and decolouration.
- 3. Classification of methods of disinfection of water.
- 4. Advantages and disadvantages of different methods of water disinfection.

Notes and problems solving

	1	r	r	1	1	1	1	1	1	r									r										
																													—
																													<u> </u>
<u> </u>																													
																													
<u> </u>																													<u> </u>
-	-			-	-	-	-	-	-		\vdash		<u> </u>	-	-		-	\vdash		\vdash	-	\vdash	-	-	<u> </u>	-		\vdash	⊢
<u> </u>	-							<u> </u>																				—_'	<u> </u>
																													<u> </u>
-																													
<u> </u>																													
	_																												<u> </u>
																													
-																													
<u> </u>	-																												┣──
	_			<u> </u>	<u> </u>		<u> </u>																					\vdash	<u> </u>
<u> </u>						-			-																				-
-	-		-	-	-	-	-	\vdash	-	-																		\vdash	
<u> </u>	-			-	-		-																						
	-			-	-		-																						
	_																											<u> </u>	<u> </u>
L								L				L	L	L	L		L				L		L	L	L	L	L	Ľ	L

Student's signature_____

Teacher's signature_____

HYGIENIC ASSESSMENT OF DRINKING WATER QUALITY. FINAL CLASS.

CONTROL QUESTIONS

- 1. Physiological, hygienic and epidemiological importance of water.
- 2. Sources of drinking water.
- 3. Water supply systems.
- 4. Methods to improve water quality (water treatment).

EXAMPLES OF CONTROL CASE TASKS

The results of laboratory investigations of water samples taken from well located near a farm

- chromaticity, degrees 40, a yellowish color;
- transparency cm 22;
- smell, points 3, swampy;
- taste, the scores 4, earthy;
- precipitate a slight, sandy;
- nitrate nitrogen (NO₃⁻), mg/l 60;
- sulphates, mg/l 150;
- chlorides, mg/l 45;
- total hardness, mEq/L 2.0;
- oxidability, mg $O_2/l 16,0;$
- the total bacterial count in 1 ml 1200.

1. Specify normative value for each of parameter. Evaluate each parameter has according to the standard value.

- 2. Make a written report on the suitability of water for drinking purposes.
- 3. What laboratory indicators suggest about water pollution with organic substances?

HYGIENIC ASSESSMENT OF SOIL

Date_____

Learning objectives:

1. To learn the chemical and physical properties of soil.

2. To learn the importance of soil pollution.

3. To learn health effects and sources of soil pollution.

4. To learn hygienic protection of the soil.

THEORETICAL SECTION

Give a definition

Soil — _____

Fill the table

Soil pollution

Pollution	Examples	Health effects	Source
Heavy metals			
Policyclic aromatic hydrocarbons			
Industrial waste			
Pesticides			

Ways humans can be exposed to soil contaminants:

1.	
2.	
3.	
з. Л	
4.	

Fill the table

Negative consequences of soil pollution

The short term effects of human exposure	1. 2. 3. 4.
Long-term effects of human exposure	1.

Possible solutions to reduce soil pollution:



KEY QUESTIONS

- 1. The chemical and physical properties of soil.
- 2. Self-purification of soil.
- 3. Soil pollution: importance, health effects and sources.
- 4. Prevention of soil pollution and its negative effects on health.

Notes and problems solving



Student's signature_____

Teacher's signature_____

HYGIENE OF HEALTHCARE ORGANIZATIONS. HYGIENIC ASSESSMENT OF THE GENERAL PLAN OF THE HEALTHCARE ORGANIZATION

Date

Learning objectives:

1. To learn requirements for the planning, construction and operation of healthcare organizations.

2 To learn hygienic requirements for the territory allocated for the construction of a healthcare organization.

3. To learn hygienic assessment of the general plan of the healthcare organization.

THEORETICAL SECTION

Fill the table

Types of hospital building construction

Туре	Characteristics

Zones of territory of modern hospital:

1.	
2.	
3	
<i>.</i>	

Hygienic requirements to hospital ground area choice:



KEY QUESTIONS

1. Hygienic requirements for the planning, construction and operation of healthcare organizations.

2. Ambulatory polyclinic organizations, their role in providing medical care to the population.

3. Hygienic requirements for the territory allocated for the construction of a healthcare organization: choice of location, size, zoning of the territory, landscaping.

4. Hygienic assessment of the general plan of the healthcare organization.

Notes and problems solving

Student's signature_____

Teacher's signature_____

HYGIENE OF HEALTHCARE ORGANIZATIONS. HYGIENIC REQUIREMENTS FOR THE DEVICE AND EQUIPMENT OF THE WARD SECTION OF THE HEALTHCARE ORGANIZATION DEPARTMENT

Date			

Learning objectives:

1. To learn requirements for the internal layout of a healthcare organization

2. To learn requirements for the device and equipment of the ward section of the healthcare organization department.

3. To learn hygienic aspects of the prevention of infections associated with the provision of medical care.

THEORETICAL SECTION

Structural divisions of hospital:
1
2
3
4
5
6
Give a definition
Ward section —
Disinfection —
Disinsection —
Deratization —
Nosocomial infections

The structure of hospital section includes the following rooms:



Fill the tables

Main routes of transmission of nosocomial infections

Route	Description

Prevention of nosocomial infections

Nonspecific	
Specific	

TRAINING TEST

1. Hygienic requirements to hospital ground area choice:

a. territory should be swamped and flooded by rivers, lakes, rain, melt waters;

b. territory should be placed in the settle zone of the city;

c. the area should be dry, on sandy or sabulous pure, safe soil;

d. far from sources of noise and environmental contamination, railways, airports, high-speed highways;

e. sanitary-protective zone should be 50–1000 m.

2. The proper level of subsoil waters is:

- a. 1.0 m from surface of soil;
- b. 1.5 m from surface of soil;
- c. 2.0 m from surface of soil;

3. Territory zones of a modern hospital are:

- a. industrial buildings;
- b. medical buildings;
- c. not medical buildings;

4. Requirements to the hospital site:

- a. area of green plantings should be 60 % or more;
- b. landscape gardening zone should be $25 \text{ m}^2 \text{ per } 1 \text{ bed}$;
- c. building area should be 15 % or less of total area;
- d. green plantings strip should be 15 m width into the perimeter of a hospital area;
- e. distance between medical building should be 24 m or more.

5. Systems of hospital build-up are:

a. combined;

c. separate;

b. centralized; d. decentralized;

6. Centralized system of hospital build-up:

a. all medical, medical-diagnostic and auxiliary departments of hospital are united in one building or in blocked building;

b different units are placed in separate buildings;

c. basic somatic departments are placed in main medical building;

d. infectious, maternity, children's, polyclinic, pathoanatomical departments and administrative unit are placed in separate buildings.

7. Mixed system of hospital build-up:

a. all medical, medical-diagnostic and auxiliary departments of hospital are united in one building or in blocked building;

b. different units are placed in separate buildings;

c. basic somatic departments are placed in main medical building;

d. infectious, maternity, children's, polyclinic, pathoanatomical departments and administrative unit are placed in separate buildings.

8. Benefits of decentralized build-up system:

a. all departments are well isolated;

- b. prevention of hospital-acquired infection;
- c. optimum conditions for medical-protective regimen;
- d. lengthening of all communications;
- e. duplication of some premises and equipment.

9. Structural divisions of a hospital are:

a. reception;

d. production department;

b. ward units;

- e. polyclinic.
- c. medical-diagnostic unit;

- d. not less than 0.5 m from foundation base;
- e. not less than 1.3 m from foundation base.

e. mixed.

- d. residential;
- e. landscape gardening.

10. Medical-diagnostic unit of hospital is:

- a. surgery block;
- b. department of regenerative treatment;
- c. department of functional diagnostics;

11. Zones of surgery block are:

- a. sterile:
- b. zone of strict mode;
- c. zone of limited mode:

- d. anesthesiology;
- e. X-ray department.
- d. zone of general hospital regimen;
- e. infected zone.

12. A zone of strict mode of surgery block includes the following premises:

- a. surgery room;
- b. preoperative;
- c. wardrobe of personnel;
- d. equipment room;
- e. postoperative wards with post of duty medical sister.

13. A zone of limited mode of surgery block includes the following premises:

a. preoperative;

- d. premises of surgeons and anesthesiologists;
- b. premises for diagnostic researches;
- c. sterilizing;

KEY QUESTIONS

- 1. Sanitary and anti-epidemic measures in the healthcare organizations.
- 2. Organization and implementation of disinfection in healthcare organizations.

3. Hygienic aspects of the prevention of infections associated with the provision of medical care.

4. Hygienic assessment of the internal layout of a healthcare organization.

5. Hygienic requirements for the device and equipment of the ward section of the healthcare organization department.

6. Sanitary and epidemiological requirements for the treatment of medical waste.

Notes and problems solving

										· ·						

Student's signature

e. gypsum room.

HYGIENE OF CHILDREN AND ADOLESCENTS. ASSESSMENT OF PHYSICAL DEVELOPMENT OF CHILDREN AND ADOLESCENTS

	Date
l 2 adolesce	Learning objectives: 1. Studying criteria for health of children and adolescents. 2. To learn methods of research and assessment of physical development of children and ents.
	THEORETICAL SECTION
Give a a	lefinition
	Growth —
I	Development —
	Fo assess the physical development of indicators are used:
1	L
2	2
3	3
Ι	Methods of assessing the physical development of children and adolescents:
1	l
2	2
3	3

PROBLEMS SOLVING

Example. During assessing the nutritional status studied physical development of 1nd year students. The obtained data were compared with individual arithmetic surveyed group (female, 17 years, time of working, sleeping, social activity 8 hours each):

Sign	Individual data (a)	М	σ	Difference between individual data and M (a-M)	Sigma deviation (a-M)/σ
Height	165	162	±2,1	+3	+3/2.1 = +1.43
Weight	61	62	$\pm 1,8$	-1	-1/1.8 = -0.56
Chest circumference	83	84	±2,0	-1	-1/2.0 = -0.5

Draw a profile of physical development of student and assess it.



disproportional physical development.

Task. During assessing the physical development of child, the individual data need to be compared with average statistical values for the age of the child (female, 9 years):

Sign	Individual data (a)	Μ	σ
Height	131	132.52	±6.24
Weight	35.5	28.06	±4.97
Chest circumference	59	62.44	±4.62

Draw a profile of physical development of child and assess it.

TRAINING TEST

- 1. Types of physical development of children and teenagers are:
 - a. disharmonious; c. functional; e. abnormal.
 - b. harmonious (normal); d. physiological;

2. The criteria characterizing health level of children are:

- a. chronic or acute diseases;
- b. level and degree of mental and physical development harmony;
- c. functional state of organism systems;
- d. degree of organism's resistance;
- e. presence or absence of allergic reactions.

3. The first place of children and adolescents' chronic diseases occurs:

- a. neuropsychiatric disorders;
- c. nasal disease (chronic tonsillitis);
- b. incorrect posture, flat feet;
- d. short-sighted.

4. The acceleration reasons are:

- a. increase in infectious diseases;
- b. influence of a city way of life;
- c. nutrition is more balanced;
- d. improving health care;
- e. endogenic.

5. Laws of growth and development of children are:

- a. dependence of growth and development processes on age;
- b. development of the musculoskeletal system;
- c. tripling of body weight at 12 months;
- d. non-uniformity of growth and development processes.

KEY QUESTIONS

- 1. Hygiene of children and adolescents: definition, purpose, objectives.
- 2. Biological and chronological age of a child. Criteria for assessment of biological age.
- 3. Criteria for health of children and adolescents. Health groups.
- 4. Methods of research and assessment of physical development of children and adolescents.

Notes and problems solving

												· · ·			· · ·		 			<u> </u>		 	 		 	 		
<u> </u>																						 					\vdash	
<u> </u>																	 					 						
<u> </u>																	 					 	 					
<u> </u>																						 					\vdash	
																	 					 	 				<u> </u>	
<u> </u>																	 				\vdash							
<u> </u>	-			<u> </u>				-	-		<u> </u>	<u> </u>	<u> </u>	<u> </u>			 		<u> </u>	<u> </u>				<u> </u>	<u> </u>		\vdash	<u> </u>
<u> </u>																						 					\vdash	
																											\vdash	
<u> </u>																												
<u> </u>								-	-												\vdash						\vdash	
<u> </u>								-									 				\square						\vdash	-
																	 					 	 					<u> </u>
]					ן ן		
1	1	1	1		1	1	1	1	1	1					1	1		1									1	(

Student's signature_____

Teacher's signature_____

FINAL CLASS ON TOPICS «THE STRUCTURE OF THE HUMAN ENVIRONMENT AND IT'S HYGIENIC SIGNIFICANCE», «HYGIENE OF HEALTHCARE ORGANIZATIONS», «HYGIENE OF CHILDREN AND ADOLESCENTS»

THE STRUCTURE OF THE HUMAN ENVIRONMENT AND IT'S HYGIENIC SIGNIFICANCE. CONTROL QUESTIONS

1. Microclimate. Optimal, acceptable microclimatic conditions. Hygienic requirements for the microclimate of residential premises and public buildings.

2. Physical properties of air (temperature, humidity): research methods, hygienic assessment.

3. Physical properties of air (barometric pressure, speed of air movement): research methods, hygienic assessment. Wind rose.

4. Hygienic assessment of combined impact of meteorological factors on the human body: methods and their characteristics.

5. Heat exchange with the environment. Consequences of its violations: overheating, overcooling. Preventive measures.

6. Chemical composition of the air environment, its hygienic characteristics. Hygienic characteristics of natural and anthropogenic sources of air pollution, their impact on health.

7. Hygienic significance of solar radiation. Pathological conditions associated with insufficient or excess solar radiation. Prevention of ultraviolet deficiency.

8. Climate and weather. Acclimatization. Meteotropic diseases and their prevention.

9. Physiological and hygienic significance of water. Water sources for drinking water supply.

10. Hygienic requirements to the quality of drinking water. Evaluation criteria.

11. The epidemic value of water. Infectious diseases transmitted by water. Methods of water purification and disinfection.

12. The chemical and physical properties of soil. Self-purification of soil.

13. Natural lighting of dwellings: hygienic significance, criteria and evaluation methods.

14. Artificial lighting: hygienic significance, research methods. Comparative hygienic assessment of various sources of artificial lighting.

15. Indoor air environment. Sources of air pollution of residential and public premises. Indicators of indoor air cleanliness.

16. Ventilation, its hygienic value. Types of ventilation and their characteristics. Indicators of the effectiveness of ventilation of premises.

17. Sources of air pollution and hygienic requirements for ventilation in the premises of healthcare organizations. Criteria for the cleanliness of the air of the premises of healthcare organizations. The role of ventilation in the prevention of nosocomial infection.

HYGIENE OF HEALTHCARE ORGANIZATIONS. CONTROL QUESTIONS

1. Hygienic requirements for the hospital area and its layout in the construction of hospitals. Functional areas allocated on the territory of health care organizations.

2. Types of hospital building construction and their hygienic assessment.

3. Requirements for internal planning of healthcare institutions. Ward section.

4. Hygienic requirements for planning and sanitary-hygienic regime maternity hospital, obstetric departments of hospitals.

5. Hygienic requirements for design and arrangement of infectious disease hospitals. Prevention of nosocomial infections

HYGIENE OF CHILDREN AND ADOLESCENTS. CONTROL QUESTIONS

1. Hygiene of children and adolescents: definition, purpose, objectives.

- 2. Biological and chronological age of a child. Criteria for assessment of biological age.
- 3. Criteria for health of children and adolescents. Health groups.

4. Methods of research and assessment of physical development of children and adolescents.

HYGIENIC ASSESSMENT OF ENERGY VALUE AND NUTRITIONAL ADEQUACY OF THE DIET. CRITERIA FOR RATIONING THE BODY'S NEEDS FOR NUTRITION, PHYSIOLOGICAL REQUIREMENTS FOR THE NUTRITION OF THE POPULATION. METHODS FOR DETERMINING THE BODY'S NEEDS FOR ENERGY AND NUTRIENTS

Date_____

Learning objectives:

1. Studying methods of determining the daily energy consumption and the needs for basic nutrients in accordance with the physiological norms of nutrition.

2. Studying the theoretical concepts of quantitative and qualitative characteristics of rational nutrition.

THEORETICAL SECTION

Foods are classified according to their functions under the following heads:

1.	
2.	
3.	

Give a definition

Total energy expenditure (TEE) — _____

Basal metabolic rate (BMR) — _____

Physical activity level (PAL) — _____

Methods for determining energy expenditure:

2	
2	
3	
4	

Fill the table

Rational or adequately nutrition

Macronutrients	Functions	Food sources
Proteins		
Fats		
Carbohydrates		

Laws of rational (adequate) nutrition:



Protein requirement is ____ g/Mcal (____ % animal origin), fat requirement is ____ g /Mcal (____ % plant origin), carbohydrates requirement is ____ g /Mcal.

PRACTICAL WORK

Calculation of individual daily total energy expenditure using physical activity level

1. Find the value of basal metabolic rate (BMR) in the table «Basal metabolic rate (BMR, kcal/day)».

2. Calculate value of BMR per 1 h. (BMR/h) = BMR/24

3. Calculate the duration of the various activities (sleep, study or work, housework, passive rest) per day

4. Calculate your energy expenses for each type of activity using the physical activity level (PAL, table «Physical activity levels for various types of activities»).

The results of calculations should be made into the table:

 $TEE = EE_{sleep} + EE_{study/work} + EE_{high active time} + EE_{low active time}$

 $E_1 = BMR/h \times Spending time_1 \times PAL_1$

Activity	Time, h	BMR/h, kcal/hour	PAL	Energy expenditure, kcal
1. Sleep and rest in bed				
2. Studying (working)				
3. High activity time (dancing, washing, ironing etc.)				
4. Low activity time (reading, watching TV, etc.)				
Total	24 h			

BMR = ____; BMR/h = ____.

Calculation of individual physiological needs in basic nutrients and biologically active substances

Using the calculated TEE and the tables «The balanced Megacalorie» and «Norms of physiological requirement for nutrients and energy for men and women», fill in the Recommended column in the protocol on the pages 61–62.

PROBLEMS SOLVING

Example. Calculate total energy expenditure of the student (male, age — 22, 70 kg; studies 8 hours, sleeping — 8 hours, social activity — 4 hours, resting — 4 hours).

- 1. BMR = 1750 kcal/day
- 2. BMR/h = 1750/24 = 73 kcal/hour
- 3. TEE = 584 + 818 + 876 + 438 = 2716 kcal

Activity	Time h	РАТ	BMR/h,	Energy expenditure,
Activity	1 1110, 11	IAL	kcal/hour	kcal
1. Sleep and rest in bed	8	1	73	$8 \times 1 \times 73 = 584$
2. Studying	8	1.4	73	$8 \times 1.4 \times 73 = 818$
3. High activity time	4	3	73	$4 \times 3 \times 73 = 876$
4. Low activity time	4	1.5	73	$4 \times 1.5 \times 73 = 438$
Total	24 h			2716 kcal

Example. The total energy expenditure of the teacher is 1920 kcal. Calculate proteins, fats and carbohydrates physiological requirements by using the balanced Megacalorie.

- 1. 1920 kcal = 1.92 Mcal
- 2. Proteins requirement = 1.92×30 g/Mcal = 57.6 g
- 3. Animal origin proteins requirement = $57.6 \times 0.6 = 34.6$ g
- 4. Fats requirement = 1.92×33 g/Mcal = 63.4 g
- 5. Plant origin fats requirement = $63.4 \times 0.3 = 19$ g
- 6. Carbohydrates requirement = 1.92×145 g/Mcal = 278.4 g

Task. Calculate total energy expenditure for teacher (male, 45 y.o., 75 kg; sleeps 8 hours, works 8 hours, low activity is 5 hours, high activity is 3 hours) using physical activity levels. Also find proteins, fats and carbohydrates physiological requirements by using balanced Megacalorie.

KEY QUESTIONS

1. Rational (adequate) nutrition. The laws of rational (adequate) nutrition.

2. Methods for determination of energy expenditure.

3. The law of nutrient (plastic) adequacy of nutrition. Formula of balanced diet.

4. The proteins and their role in human nutrition. Foods that are the main sources of complete protein.

5. Fats and their role in human nutrition. Foods that are major sources of fats.

6. Carbohydrates and their role in human nutrition. Foods that are major sources of carbohydrates.



Notes and problems solving

Student's signature_____

Teacher's signature_____

HYGIENIC ASSESSMENT OF THE ENERGY VALUE AND NUTRITIONAL ADEQUACY OF THE DIET. CALCULATION OF THE ACTUAL NUTRIENT INTAKE AND ENERGY VALUE OF THE DIET ACCORDING TO THE 24-HOUR RE-CALL METHOD

Date

Learning objectives:

1. Studying methods of determining the actual nutrient intake and energy value of the diet.

2. Studying the theoretical concepts of quantitative and qualitative characteristics of rational nutrition.

THEORETICAL SECTION

The methods are most commonly used to assess food intake:

1.	
2.	
3.	
Δ	
••	

Protein provides _____ calories/g, fat provides _____ calories/g, carbohydrates provide _____ calories/g (inscribe the correct numbers).

PRACTICAL WORK

Case study «24-hours dietary re-call»

Estimate the actual nutrients intake using the 24-hours dietary re-call method. Fill out the table on page 56–57 and calculate the actual intake of nutrients. Put the result in the Actual column in the protocol on the pages 58–59.

PROBLEMS SOLVING

Task. Student (age -22, body weight -76 kg; studying for 7 hours, sleeping -8 hours, social activity -5 hours, resting -4 hours) intakes 74 g of proteins, 87 g of fats and 319 g of carbohydrates. Calculate total energy intake and compare it with the total energy expenditure.

KEY QUESTIONS

1. Rational (adequate) nutrition. The laws (principles) of rational (adequate) nutrition.

2. Methods of determining the actual consumption of nutrients and energy.

Case study «24-hours dietary re-call»

Food consumption data:

Food							Intak	æ			
Food	Amount, g	protein, g	fat, g	carbs, g	Ca	Fe	Mg	Vitamin C	Vitamin A	Vitamin D	Energy
Breakfast											
Lunch											
Dinner											

59

End of the table

Food							Intal	ĸe			
Food	Amount, g	protein, g	fat, g	carbs, g	Ca	Fe	Mg	Vitamin C	Vitamin A	Vitamin D	Energy
Dinner											
Additionally ×											
Snacks											
Total											

* Brunch/afternoon snack

60

Notes and problems solving

<u> </u>					-														

Student's signature_____

Teacher's signature_____

HYGIENIC ASSESSMENT OF THE ENERGY VALUE AND NUTRITIONAL ADEQUACY OF THE DIET. ASSESSMENT OF THE ADEQUACY OF ACTUAL DIET TO THE NEEDS OF THE HUMAN BODY, RECOMMENDATIONS FOR RATIONALIZING NUTRITION

Date_____

Learning objectives:

1. Studying methods of hygienic assessment of the energy value and nutritional adequacy of the diet.

2. Studying methods of rationalizations nutrition.

PRACTICAL WORK

Hygienic assessment of the adequacy of actual nutrition

Nº	Te de se de ser	TT \$4	December 1.1	A	Devi	ation
JN⊇	Indicators	Unit	Recommended	Actual	Lack	Excess
1.	Energy value	kcal				
2.	Proteins,	g				
	animal origin proteins	g				
3.	Fats,	g				
	Plant origin fats	g				
4.	Carbohydrates	g				
5.	Minerals:					
	Са					
	Fe	mg				
	Mg					
6.	Vitamins:					
	С	-				
	А	mg				
	D					
7.	Calorific value by:					
	Proteins		12			
	Fats	%	30			
	Carbohydrates		58			

End of the table

NG	Indiastans	T	Decem	ded	A stual	Devi	ation
JN⊇	Indicators	Unit	Recom	nended	Actual	Lack	Excess
8.	Ratio (by mass) Proteins : Fats : Carbs	g	1:1.2	2:4.6			
9.	Calorific value by:		4 meals a day	3 meals a day			
	breakfast		25	30			
	lunch		35	40 - 45			
	afternoon snack dinner		15	_			
			25	25 - 30			

Conclusion: _____

Recommendations for rationalizing nutrition: — _____

PROBLEMS SOLVING

Task. Nurse (female, 45 y. o., 60 kg) sleeps 8 hours, works 8 hours, has 4 hours of low activity and 4 hours of high activity per day. Intake of nutrients is 75 g of proteins, 100 g of fats and 350 g of carbohydrates.

Calculate her total energy expenditure, requirements for macronutrients, actual energy intake. Compare her actual energy intake and total energy expenditure, actual intake of nutrients and requirements.

Make a conclusion about her diet. Is it rational? How to improve her diet?

TRAINING TEST

1. Energy body use for ingestion and digestion of food, and for the absorption, transport, interconversion, oxidation and deposition of nutrients is called:

a. basal metabolism;

- b. growth;
- c. dietary-induced thermogenesis;
- d. physical activity level.

2. The metabolic response to food increases total energy expenditure by:

a. about 78 % of the BMR over a 24-hour period in individuals with a mixed diet;

- b. about 10 % of the BMR over a 24-hour period in individuals with a mixed diet;
- c. about 1 % of the BMR over a 24-hour period in individuals with a mixed diet;
- d. about 50 % of the BMR over a 24-hour period in individuals with a mixed diet.

3. According to the law of the energetic adequacy of nutrition:

- a. food must be cooked in accordance to the digestive system enzymic abilities;
- b. the dietary intake food calorific value must correspond to the organism energy consumption;
 - c. food must not be toxic and must be harmless in epidemiological aspect;
 - d. food intake hours must correspond to the organism biological rhythms.

4. According to the law of nutrient (plastic) adequacy of nutrition:

- a. food must be cooked in accordance to the digestive system enzymic abilities;
- b. food must not contain the etiological agents of infectious foodborne diseases;
- c. food intake hours must correspond to the organism biological rhythms;
- d. all nutrients must be contained in optimal quantities and ratios.

5. Principles of rational diet:

- a. the last food intake must be not less than 2–2.5 hours before sleeping;
- b. adults must have 3-4 meals a day;
- c. intervals between food intakes must be less than 5 hours;
- d. intervals between food intakes must be more than 5 hours in day time;
- e. adults must have 2–3 meals a day.

6. The group of energy-yielding foods include:

- a. foods rich in protein;
- b. foods rich in carbohydrate and fat;
- c. foods rich in vitamins, minerals.

7. Classification of types of nutrition includes:

- a. intuitive nutrition; c. rational nutrition;
- b. preventive nutrition; d. clinical nutrition.

8. Sources of unsaturated fats are:

- a. meat products, meat pies, sausages;
- b. oily fish;
- c. cakes and biscuits;
- d. nuts and seeds.

9. According to the formula of balanced nutrition for adults:

- a. the ratio between proteins, fats, carbohydrates is 1:1:2;
- b. the ratio between proteins, fats, carbohydrates is 1 : 1.1-1.2 : 4.6;
- c. the ratio between proteins, fats, carbohydrates is 1:4:9.

10. The amount of energy used for basal metabolism in a period of time is called ______

11. According to the law of biotic adequate nutrition:

a. the dietary intake food calorific value must correspond to the organism energy consumption

- b. food must not be toxic and must be harmless in epidemiological aspect
- c. food must be cooked in accordance to the digestive system enzymic abilities

d. food intake hours must correspond to the organism biological rhythms

12. The main sourses of proteins are:

a. fruits;	e. milk;
b. vegetables;	f. meat;
c. fish;	g. eggs
d. seeds;	

13. Carbohydrate requirements:

a.	4	g/Mcal;	
----	---	---------	--

b. 33 g/Mcal;

c. 1 g/kg body weight

d. 145 g/Mcal;e. 4 kcal/1 g.

KEY QUESTIONS

1. The law of nutrient (plastic) adequacy of nutrition. Classification of nutrients. Formula of balanced diet.

2. Methods for determination of energy expenditure and methods of determining the actual consumption of nutrient energy.

3. Classification of nutrients. Formula of balanced diet.

4. The impact of nutrition on health. Preventive and therapeutic role of nutrition.

5. Hygienic assessment of adequacy of dietary requirements of an organism, the development of recommendations for optimizing nutrition.

1 1																								
1 1																								
1 1																								
I I																								
1 1																								
I I																								
1 1																								
Image: Solution of the state of the sta																								
Image: Solution of the state of the sta																								

Notes and problems solving

Student's signature

Teacher's signature

HYGIENIC ASSESSMENT OF NUTRITIONAL STATUS. DETERMINATION OF SOMATOMETRIC, SOMATOSCOPIC, PHYSIOMETRIC INDICATORS

Learning objectives:

Date_____

1. Nutritional status definition, stages of nutritional status assessment.

- 2. Classification of nutritional status.
- 3. Criteria and methods of nutritional status assessment.

THEORETICAL SECTION

Give a definition

Nutritional status is _____

Nutritional status classification										
1	1	1	1							
		2	2							
	2	3	3							
			4							

Stages of nutritional status assessment are



Methods of nutritional status assessment



Somatotype	1
Physical development, functional and adaptive reserves of the body	1 2 3
Protein, mineral, vitamin metabolism	1 2 3
Immunologicalresistance of skin and mucous membranes	1 2

PRACTICAL WORK

Methods of actual nutrition assessment

(completed on class «Hygienic assessment of the energy value and nutritional adequacy of the diet. Assessment of the adequacy of actual diet to the needs of the human body, recommendations for rationalizing nutrition»)

Methods of assessment the state of health that has developed under the influence of actual nutrition

(Put the result in the protocol on the page 68)

Assessment of somatoscopic indicators of physical development

Type of constitution	Chest shape	Epigastric angle	Leg shape ¹	Belly shape	Skeleton development (points) ²	Muscles development (points) ³	Body fat (points) ⁴
Asthenoid	Flattened	< 90°	O-shape	Sunken	1	1	1
Thoracic	Cylindrical	90° < 90°	X, H, O- shaped	Straight	1–2	1–2	1–2
Muscular	Cylindrical and conic	90° > 90°	X, H, O- shaped	Straight	2–3	2–3	1–2
Digestive	Conic	> 90 ⁰	X-shaped	Bulging	2–3	1–2	2–3

Somatoscopic characteristics of the main types of body constitution

Notes:

¹Leg shape:

O-shaped legs (genu varum), when standing with closed heels and spread toes, there is a distance between the inner edges of the knee joints, the legs are arched as if; the axis of the lower leg with the axis of the thigh forms an angle open inwards;

H-shaped legs — a straight position in which the axes of the thighs and shins are parallel, the legs are straight.

X-shaped legs (genu valgum) — the knees are shifted, the axes of the lower legs diverge, the axis of the thigh with the axis of the lower leg form an angle open to the outside.

²Development of skeleton: the massiveness of the skeleton should be noted, based on the relief and size in the area of large joints (elbow, knee, wrist). There are three grades: thin (1 point), medium (2 points), and massive (3 points).

³ Muscles development:

When evaluating musculature, it is necessary to take into account the entire musculature as a whole, and not just one segment. Evaluation of musculature is carried out in two directions: quantitative development of the muscle layer, qualitative state of the musculature (tone, elasticity).

There are three degrees of muscle development: small (1 point), medium (2 points), and large (3 points). Determination of muscle tone is established by probing the muscles in a relaxed and tense state: weak tone (1 point), medium (2 points), good (3 points). It is quite enough to give a general assessment of the musculature, taking into account both its quantitative development and the degree of elasticity (tone).

⁴Body fat deposition — development of subcutaneous adipose tissue. There are three degrees of development of fat deposition: small (1 point) — the bone relief (shoulder girdle, shoulder blades, ribs) clearly protrudes; medium (2 points) — the bone relief is not clearly expressed; large (3 points) - smoothed bone relief, rounded contours.

Conclusion: Type of constitution is

Assessment of somatometric indicators of physical development

Somatometric indicators of physical development is determined using generally accepted methods for measuring body weight and length, the circumference of the chest in pause, on inhalation and exhalation (chest excursion).

1. Standing height:

The subject stands straight, hands at the seams, heels together, toes apart. The patient touches the height meter rack with his heels, buttocks and the interscapular area. The head should be in such a position that the line mentally drawn from the upper edge of the tragus of the ear to the lower edge of the eye socket is horizontal.

2. **Body weight** measurement using medical scales: weighing is performed on an empty stomach, without clothes and shoes. The subject is standing in the middle of the weighing platform.

3. Measuring the **chest circumference** with a centimeter measuring tape. Chest circumference is measured at rest, maximum inhalation and maximum exhalation. The tape is applied from behind at the lower corners of the shoulder blades, in front along the edge of the periarticular circle (in men), in women — along the IV rib.

Body mass index	
Body mass index (BMI) = weight / (height) ² , (kg/m ²).	
Normal value of BMI is mentioned in the table.	

Category	BMI (kg/m ²)	
Underweight (Severe thinness)	< 16,0	
Underweight (Moderate thinness)	16,0–16,9	
Underweight (Mild thinness)	17,0–18,4	
Normal range	18,5–24,9	
Overweight (Pre-obese)	25,0–29,9	
Obese (Class I)	30,0–34,9	
Obese (Class II)	35,0–39,9	
Obese (Class III)	\geq 40,0	

Assessment of body composition

Measuring the thickness of the skin-fat fold using a transparent ruler or caliper at three points. 1. On the abdominal wall in the iliac region along the parasternal line (5 cm from the navel).

2. Along the mid-axial line at the level of the breast nipple.

3. At the angle of the shoulder blade.

At the same time, in the first 2 points, the skin with subcutaneous fat is captured by the fingers in the horizontal direction, at the angle of the shoulder blade — in the vertical. The measured thickness of the fat fold is divided by 2.

Actual value (average at three points) = _____

Body surface area (BSA) = $134 \times BW + 52.6 \times BH$

BW — body weight, kg; BH — standing height, cm. Actual value (body surface area) = _____

Body fat weight (BFW) = SF \times BSA \times 0.0632

SF — is the average thickness of the skin-fat fold at three points, mm; BSA — is the surface area of the body, cm^2 ; 0,0632 — is an empirical coefficient.

Actual value (body fat weight) = _____

Body fat content (BFC) = $(BFW / BW) \times 100 \%$

BFW — is the amount of body fat, g; BW — is the body weight, g. Actual value (body fat content) = _____

The amount of lean body weight (LBW) = BW — BFW

BW — is the body weight, g; BFW — is the amount of body fat content, g. Actual value (lean body weight) = _____

ACE body fat % norms				
Description	Women	Men		
Essential Fat	10–13 %	2–5 %		
Athletes	14–20 %	6–13 %		
Fitness	21–24 %	14–17 %		
Average	25–31 %	18-24 %		
Obese	32 % +	25 %+		

Assessment of phisiometric indicators of physical development

Phisiometric indicators of physical development is determined using generally accepted methods for measuring the muscular strength of the hands (by dynamometry) and the vital capacity of the lungs (by spirometry).

Measuring the muscular strength of the hands with a manual dynamometer

It is necessary to squeeze the dynamometer as much as possible with the hand extended and withdrawn to the side. Take into account the maximum result.

Strength index = (Muscle strength of the right hand (kg) / body weight (kg)) \times 100 % Normally strength index for male is 70–75 %, for females 35–40 %.

Measurement of the vital capacity of the lungs using a spirometer

The patient takes a maximum breath, holding his breath, tightly wrap his mouth around the mouthpiece and exhale all the air into the tube. The maximum indicator of 2-3 measurements is taken into account.

Vital index

Vital index = lung vital capacity (ml) / body weight (kg) Normally vital index for male is > 60 ml/kg, for females is > 50 mg/kg.

Cardio-respiratory index (CRI)

Cardio-respiratory index (CRI) = heart rate / breathing rate

At rest, the CRI is close to 4–5 and shows the consistency of the respiratory and cardiovascular systems and their interdependence. The sharp increase in the CRI indicates cardiovascular system stress, while its decrease indicates the initial signs of decompensation in the respiratory system.

Test Ghencea. Index of resistance to hypoxia

Test Ghencea with arbitrary breath holding (breath holding on exhalation) gives an indication of the reserves of oxygen for metabolic needs of the body.

At rest, the pulse is counted for 30 sec. The patient makes 3 deep breaths, then inhale — exhale and holds his breath, holding nose with his fingers. Delay time breathing recorded stopwatch. After the resumption of breathing count pulse for 30 sec.

Assessment is based on the index of resistance to hypoxia (IRH) with the breath-holding time.

Index of resistance to hypoxia (IRH) = $\frac{\text{heart rate for 30 seconds after holding your breath}}{\text{time breath holding}}$

Duration breath-holding time is evaluated by a four-point system: excellent condition — more than 50 sec; good — 30-50 sec; average — 20-29 sec; poor — less than 20 sec.

= _____ =

Normally IRH less or equal to 1 (it is less than the higher resistance to hypoxia).

Assessment of functional parameters of nutritional status. Index of physical status

The development of physical condition can be evaluated on an index of physical status (IPS).

$$IPS = \frac{(700 - 3 \times heart \, rate - \, 0.8333 \times SBP - \, 1.6667 \times DBP - \, 2.7 \times Age + \, 0.28 \times BW)}{(350 - 2.6 \times Age \, + \, 0.21 \times BH)} =$$

$$IPS = \frac{(700 - 3 \times _ - 0.8333 \times _ - 1.6667 \times _ - 2.7 \times _ + 0.28 \times _)}{(350 - 2.6 \times + 0.21 \times)}$$

where, IPS — index of physical condition; heart rate — pulse rate, beats/min; Age — calendar age, years; SBP — systolic blood pressure, mmHg; DBP — diastolic blood pressure, mmHg; BW — body weight, kg; BH — body height, cm.

IPS assessed taking into account received numerical grades for the following levels: low — less than 0,375; below average — 0,375–0,525; average — 0,526–0,675; above average — 0,676–0,825; high from — 0,826 or more.

Adaptive reserves. Adaptation potential. In order to assess the level of an integrated health and detect possible violations of adaptation is calculated integral measure of the adaptation reserves of the circulatory system — adaptation potential.

where, AP — adaptation potential; heart rate — pulse rate, beats / min; Age — calendar age, years; SBP — systolic blood pressure, mmHg; DBP — diastolic blood pressure, mmHg; BW — body weight, kg; BH — body height, cm.

Assessment of the results: satisfactory adaptation: 2.20 or less - for young men; 1.96 or less — for young women; stress adaptation: 2.21–2.43 — for young men; 1,97–2,23 — for young women; poor adaptation and failure of adaptation: 2.44 or more — for young men; 2.24 or more for young women.

TRAINING TEST

1. Correlate nutritional status and definition the number and the letter):

1. The optimal (ideal) status;

2. The normal (usual) nutritional status:

3. The insufficient status:

a) characterized by strong health and wide range of adaptive reserves;

b) charactarised by narrow range of adaptive reserves, decreased working capacity and health condition;

c) characterized by absence of functional and structural changes and adequate range of adaptive reserves.

Answer: ____

2. Normal value of BMI is:

a. 18.5–29.9; b. 19.5–25; c. 18.5–24.9; d. 22–30.

3. For the characteristics of physical development, functional and adaptive reserves of the body can be assessed using the following methods:

a. physiometric; b. somatometric; c. biochemical; d. physiological.

4. Organize chronologically the steps of nutritional status assessment from the beginning:

_____. correction of nutritional status;

_____. assessment of actual nutrition and identifying limiting factors in the diet;

_____. health assessment related to nutrition;

_____. definition of a type of nutritional status.

KEY QUESTIONS

1. Nutritional status definition, stages of nutritional status assessment.

2. Classification of nutritional status.

3. Criteria and methods of nutritional status assessment.

4. Characteristic of the common and optimal (ideal) nutrition status.

Student's signature_____

Teacher's signature

HYGIENIC ASSESSMENT OF NUTRITIONAL STATUS. RECOMMENDATIONS FOR OPTIMIZING NUTRITIONAL STATUS

Date_____

Sign	Indicator	Physiological norm	Actual value	Deviation
Intake of nutrients	1. Energy value, kcal			
	2. Proteins g/day			
	3. Fats, g/day			
	4. Carbohydrates, g/day			
Body composition	Body weight, kg			
	Body height, cm			
	Body mass index, kg/m ²	18,5–25,0		
	Body fat, %	M		
	Index of physical state units	Г Лього 0 526		
		ADOVE 0,520		
	Strength index, %	M 70 - 73 E 25 40		
		r 55–40		
	Lungs vital capacity, ml	According		
Functional		$\frac{10110gramm}{M > 60}$		
parameters Adaptation parameters	Vital index, ml/kg	M > 00, E > 50		
	Cardio respiratory index units	1 > 50		
	Cardio-respiratory index, units	<u>4–3</u> <u>30</u>		
	Duration of breath-holding, sec	30 and more		
	Index of registeries to hyperic			
	A doptation potential units	≥ 1		
	Adaptation potential, units	$\leq 2,0$		
	Morbidity (number of acute	Less than 4 cases		
	respiratory infection case within	in a year		
	previous year)	-		

Nutritional status assessment protocol

Conclusion _____

Type of nutritional status _____

Recommendations for optimizing nutritional status

Note: for the stabilization of body weight in the case of excess or lack is necessary to increase (reduce) the energy value of the daily diet of the calculated value. Weight gain of 1 kg correspond to excess consumption of food calories 6800 kcal. Therefore, the correction of the daily diet for a month will be 6800 kcal/30 day = 227 kcal/day.
PROBLEMS SOLVING

Task. During assessing the nutritional status studied physical development of students. The obtained data were compared with individual arithmetic surveyed group (male, 20 years, time of working, sleeping, social activity 8 hours each).

Sign	Individual data (a)	М	σ	Difference between individual data and M (a-M)	Sigma deviation (a-M)/σ
Height	181	176	±2.3		
Weight	72	74	±2.1		
Chest circumference	94	93	±2.4		

Draw a profile of physical development of student and assess it.

KEY QUESTIONS

1. Nutritional status definition, stages of nutritional status assessment.

2. Classification of nutritional status.

3. Criteria and methods of nutritional status assessment.

4. Characteristic of the common and optimal (ideal) nutrition status.

5. Alimentary diseases: definition of the concept, causes, classification.

6. Insufficient nutrition status. Hypotrophy, alimentary dystrophy and dwarfism: clinical symptoms, prevention.

7. Diseases of protein-energy malnutrition: classification, diagnosis, clinic, prevention.

8. Excess nutrition status. Obesity: causes, classification, diagnosis, prevention.

9. Protein excess nutrition syndrome: causes of development, manifestations, prevention.

Notes and problems solving

Student's signature_____

Teacher's signature_____

HYGIENIC ASSESSMENT OF THE PROVISION OF THE HUMAN BODY WITH VITAMINS. RECOMMENDATIONS FOR OPTIMIZING THE VITAMIN SUPPLY OF THE HUMAN BODY

Date_____

Learning objectives:

1. Classification and physiological significance of vitamins in the organism.

2. To master the methods of detection and assessment of the vitamin sufficiency in the organism and the methods and measures of hypo- and avitaminosis prevention.

3. The most frequently occuring hypovitaminosis states in cases of both individual and collective nutrition. Their causes.

4. Avitaminosis and their clinical characteristics.

5. Hygienic principles of prophylaxis of hypovitaminosis and avitaminosis.

6. Detecting the hypo- and avitaminosis in cases of both individual and collective nutrition.

7. Organizing the hypovitaminosis prevention measures and assess their effectiveness.

8. Revealing the deficit of vitamins in the human organism by determining of the quantity of vitamin C in urine and to estimate the results.

THEORETICAL SECTION

Give a definition Vitamins — _____ Fill the table **Classification of Vitamins** Vitamins **Types of Vitamin Status of Organism** By level of vitamin supply of organism the following are distinguished: Avitaminosis — Hypovitaminosis — _____ Normal supply — _____ Hypervitaminosis — _____

Fill the table

Vitamin	Sources	Functions of Vitamin	Symptoms of Vitamin Deficiency	Daily Requirement of Vitamin
vitamin A				
vitamin D				
vitamin E				
vitamin V				
vitainin K				
vitamin B ₁				
vitamin B ₂				
vitamin B ₃				
vitamin B ₅				
vitamin B ₆				

End of the table

Vitamin	Sources	Functions of Vitamin	Symptoms of Vitamin Deficiency	Daily Requirement of Vitamin
vitamin B7				
vitamin B9				
vitamin B ₁₂				
vitamin C				

Methods of diagnostics of vitamin status are as follows:

The study of dietary vitamin and actual consumption of vitamins from food	The study of the vitamin status of the organism.

PROBLEMS SOLVING

Example. The daily diet of patient contains 180 g of cabbage. The content of vitamin C in cabbage is 50 mg per 100 g. During cooking average vitamin C losses is 60 %. There is a decrease of skin capillary resistance. Excretion of ascorbic acid in the urine was 0.3 mg/h.

1. amount of ascorbic acid intake (excluding losses):

50 mg % - 100g

X mg % – 180g,

 $X = 180 \times 50/100 = 90 \text{ mg }\%$

2. actual intake (taking into account cooking loss of vitamin C):

90 mg - 100 %

Y mg - 40 %

 $Y = 40 \times 90/100 = 36 \text{ mg} / \text{day}$

3. estimate the amount of vitamin C that is excreted in the urine:

You must remember that with sufficient intake of vitamin C, its excretion in the urine is 0.7-1.0 mg/day.

In this task, excretion of as corbic acid in the urine was 0.3 mg/day — indicating a lack of vitamin $\rm C$

4. clinical manifestations must be evaluated:

In this task, a decrease of skin capillary resistance indicating a lack of vitamin C

Conclusion: insufficient amount of vitamin + nutritional recommendations (it is necessary to list the products — sources of vitamin C)

Tasks. 1. Patient complained a feeling of weakness and pain in the legs, leg fatigue when walking. In an interview with the patient have been revealed significant errors in its food: high daily consumption of confectionery products, regular use in nutrition of wheat bread flour, cream of wheat and porridge cooked from polished rice. During palpation found pain calves. Excretion of ascorbic acid in the urine was 0.2 mg/h. There is an increase of capillary fragility of the skin. What vitamins defficiency cause similar symptoms?

2. Patient complained of increased frequency of respiratory diseases and severe worsening ability to see objects at dusk. It is revealed that the patient is a strict vegetarian. Examination revealed follicular hyperkeratosis and disruption of normal color vision. What vitamins defficiency cause similar symptoms?

3. Student (male, 21 y.o., 75 kg) sleeps 7 h, studies 9 h, has 5 h of low activity and of 3 h high activity. The daily diet contains 50 g of broccoli as a vitamin C source. The content of vitamin C in broccoli is 89 mg per 100 g. During cooking average, vitamin C loss is 65 %. Objective data: increase in capillary fragility of the skin, decreased productivity. Excretion of ascorbic acid in the urine was 0.2 mg/h.

Assess providing the body with vitamin C.

TRAINING TEST

1. Vitamins — are essential substances that:

a. have plastic value;	d. perform catalytic functions;
b. have the energy value;	e. contained in drinking water.
c. are macronutrients;	
2. Water-soluble vitamins include:	
a. beta-carotene;	c. vitamin E (tocopherol);
b. folic acid;	d. vitamin K.
3. Fat-soluble vitamins include:	
a. vitamin K;	d. vitamin C;
b. pantothenic acid;	e. vitamin B_{12} .
c. choline:	

4. Specify the most characteristic symptom of vitamin A deficiency:

a. skin cyanosis;

b. swelling interdental papillae;

c. hypothermia;

d. formation of petechiae;

e. increase in the duration phase of dark adaptation.

5. Specify the most characteristic symptom of vitamin C:

- a. follicular hyperkeratosis (goosebumps) in combination with petechiae;
- b. burning sensation in the feet;

c. conjunctivitis and blepharitis;

- d. corneal vascularization;
- e. night blindness.

6. Select a product in which the vitamin C content greater than 100 mg %: e. sunflower oil.

- c. fish; a. liver;
- d. rosehip; b. eggs;

7. A normal amount of ascorbic acid excreted in the urine is:

a. 0.5–0.6 mg / hour;	c. 0.7–1.0 mg / hour;
b. <0.2 mg / hour;	d. 0.03–0.25 mg / hour.

8. The main vitamin A provider:

a.cod liver;	c. potatoes;	e. citrus.
b. bran;	d. cabbage;	
Insufficiency of which w	itamin aan aawaa nallaan	

9. Insufficiency of which vitamin can cause pellagra:

a. folic acid; c. niacine: e. cyanocobalamin. b. pantothenic acid; d. biotine;

10. Select a product in which the vitamin C content is above 100 mg %:

a. cauliflower; c. strawberry; e. garnet.

b. citrus; d. red bell pepper;

KEY QUESTIONS

- 1. The physiological role of vitamins and hygienic norms in daily ration.
- 2. Main food sources of vitamin C in nutrition.
- 3. Etiology, pathogenesis, symptoms of avitaminosis and hypovitaminosis C.
- 4. The methods of vitamin C sufficiency control in the human organism.
- 5. The method of vitamin C determination in urine (by Thilman's reagent).
- 6. Vitamins: definition, classification, functions.
- 7. Methods of assessing supply the body with vitamins.
- 8. Vitamin A: physiological significance, the daily demand, the main sources.
- 9. Vitamin C: The physiological significance of the daily demand, the main sources.

						 	 . 1	 	 	-	0						

Notes and problems solving

Notes and problems solving

																						 '	<u> </u>
<u> </u>																							
										 							 		 				<u> </u>
<u> </u>																						\vdash	
																							<u> </u>
<u> </u>										 							 		 			 \vdash	
																							<u> </u>
<u> </u>										 							 		 	 		 	
<u> </u>							-															\vdash	<u> </u>
																							
<u> </u>																							
																						\square	
																							<u> </u>
<u> </u>										 							 		 				-
																							<u> </u>
<u> </u>																							
<u> </u>																						\vdash	
																							<u> </u>
<u> </u>																							
<u> </u>							-													 		 \vdash	┣──
<u> </u>																							<u> </u>
																							1
<u> </u>		 -					-		-		-	 		-	 			-		 -	-	 \vdash	
<u> </u>		 										 			 					 		 ├ ──	├
																						\vdash	<u> </u>
<u> </u>		 										 			 					 		 ┝──╵	├
<u> </u>				<u> </u>		<u> </u>																<u> </u>	<u> </u>

Student's signature_____

Teacher's signature_____

PREVENTION OF FOOD POISONING OF MICROBIAL ETIOLOGY

	Date
	Learning objectives:
	1. Studying the law of biotic adequacy of nutrition.
	2. Food poisoning of microbial nature: toxiconflections. Preventive measures.
	4. Food poisoning of microbial nature: mycotoxicoses and phycotoxicoses. Preventive
mea	sures.
	THEORETICAL SECTION
Give	a definition
	Food poisoning —
	Food poisoning classification:
1.	
_	a.
	D
2	
	a
	b.
3	C
5	
	Triple condition for the occurrence of foodborne diseases (infections):
	1
	2
	3
	3
	Additional factors for the occurrence of foodborne diseases (infections):
	1
	2.

Fill the table

Nosological form		Etiological factor
Toxico infection	•	
	•	
	•	
	•	
	•	
	•	
	•	
	•	
Toxicosis: •	•	
•	•	
•	•	
Mixed	•	
	•	

The epidemiological pattern of foodborne disease (infection) manifestations:

4	4
5	5
e	5
7	7
8	3
S	Sources of food products contaminations with S. aureus:
S	Symptoms of botulism:
I	Preventive measures of bacterial food poisoning:

PROBLEMS SOLVING

Task. In the village, 3 families applied for medical help, presenting the following complaints: abdominal pain, repeated profuse vomiting, weakness, dizziness, cold sweat. 8 people got sick. All the families consumed various foods on the day of the disease. Common to all the products was milk, which was systematically bought from a resident of a neighboring village. On the day before the disease, in all cases, milk was consumed without pre-boiling (it was added to coffee, cocoa, to dilute porridge).

The day before the disease, a woman selling milk warned customers about stop delivering milk due to a veterinarian warning about the need to treat the cow for mastitis.

Determine the presumptive diagnosis and suspected product. Which priority measures must be taken?

Fill the table

Mycotoxicoses

Mycotoxins	Main producers	Sources of mycotoxins	Toxic effect
Aflatoxins B1, M1			
Trichothecene mycotoxins:			
dezoxinivalenol			
T-2 toxin			
Patulin			
Ochratoxin A			
Fumonisin B1			
Ergoalkaloids			

Phycotoxicoses

Phycotoxins	Main producers	Sources phycotoxins	Toxic effect
Group PSP (paralytic shell- fish poison): saxitoxin			
Group DSP (diarrheic shell-fish poison):okadaic acid, pektenotoxin			
Group neurotoxic: Brevetoxins			
Amnestic group: domoic acid			

KEY QUESTIONS

1. The law of biotic adequacy of nutrition.

2. Food poisoning: definition, classification.

3. Food poisoning of microbial nature: toxicoinfections. Preventive measures.

4. Food poisoning of microbial nature: bacteriotoxicosis. Preventive measures.

5. Food poisoning of microbial nature: mycotoxicoses and phycotoxicoses. Preventive measures.

Notes and problems solving

								1																		
																										<u> </u>
									-																\vdash	
<u> </u>																									<u> </u>	<u> </u>
																										
<u> </u>											 														\vdash	
																										<u> </u>
<u> </u>											 				 				 		 					-
											 				 				 		 					<u> </u>
<u> </u>											 				 				 		 					
<u> </u>											 				 				 		 				 \vdash	
																										<u> </u>
-																										
<u> </u>																										
<u> </u>																										_
																										1
<u> </u>															 											
<u> </u>																									—	-
<u> </u>																										
<u> </u>			-					-	-	-		 	-	\vdash		\vdash	 	\vdash		-		-	-	-	 \vdash	⊢
<u> </u>												 			 		 								 ├ ─-	_

Student's signature_____

Teacher's signature_____

PREVENTION OF NON-MICROBIAL FOOD POISONING

Learning objectives:

1. Studying food poisoning of non-microbial nature classification.

2. Studying poisoning with chemical impurities and their prevention.

3. Studying food poisoning with products of plant and animal origin: products that are poisonous in nature and products, poisonous under certain conditions, their prevention.

4. Studying food poisoning with poisonous mushrooms and plants prevention.

5. Studying food poisoning detecting.

THEORETICAL SECTION

Fill the table

	Poisoning with products that are poisonous in nature
Plant origin	Wild and cultivated plants
	1
	2
	3
	4
	5
	0
	8
	9
	10.
	Seeds of weeds of cereal crops
	1
	2
	3
	4
	5
	Poisonous mushrooms
	1
	2
	3
Animal	Internal organs of individual fish species
origin	Internal organs of individual fish species
ongin	1
	2
	3
	4
	7
	7:
	Some endocrine glands of slaughter animals
	1.
	2

Date_____

End of the table

	Poisoning with poisonous products under certain conditions
Plant origin	Kernels of fruit seeds 1.
	2
	nuts;
	beans, containing;
	potato, containing
	Conditionally edible mushrooms
	1.
	4, not subjected to proper culinary processing.
	Edible mushrooms —
Plant origin	Liver, caviar of some fish species during the spawning period 1.
	Mussels and crustaceans acquiring toxic properties
	Honey
Poisoning with impurities of toxic substances xenobiotics	1.

Preventive measures of non- microbial food poisoning:

What should a doctor do if a patient is suspected of food poisoning?

PROBLEMS SOLVING

Task. Five of younger schoolboys ate broken apricots kernel. After 5 hours, the two children appeared dizziness, headache, nausea, general weakness (which have passed within 6 hours). One boy after 3 hours started vomiting, severe headache, dizziness, and diarrhea. The two children who have eaten most kernels seed, in 2 hours all the above mentioned symptoms were strongly expressed, one fainted. On examination, the doctor revealed a cyanotic face and mucosal conical and tonic spasms, trismus. The next day the one child died. In other children, after intensive therapy, the condition improved and they have recovered.

Determine the presumptive diagnosis and suspected product. Which priority measures must be taken?

TRAINING TEST

1. Food poisoning are:

a. usually chronic diseases;

b. rarely chronic diseases;

2. Classification of food poisoning includes:

- a. microbial food poisonings;
- b. viral food poisonings;
- c. food poisonings caused by plant;
- 3. Classification of microbial food poisoning includes:
 - a. foodborne diseases; b. food poisoning unknown etiology;
- d. food toxicosis;
- e. mixed food poisoning.
- c. food poisoning caused by chemical substances;

4. Classification of non-microbial food poisoning includes:

- a. poisoning caused by fungi toxins;
- b. poisoning caused by endocrine glands of animals (adrenal, pancreas);
- c. poisoning caused by canned food containing botulinum toxin;
- d. phycotoxycosis.

c. rarely acute disease;

d. usually acute disease.

- e. non-microbial food poisonings.
- d. unknown etiology;

5. Doctor who diagnosed or suspecting food poisoning, has to:

a. notify about food poisoning Ministry of healthcare;

b. immediately transport the patient to the infectious department an ambulance;

c. notify about food poisoning Center of Hygiene and Epidemiology;

d. immediately prohibit further eating of suspected food;

e. take samples of suspected food, collect vomit, feces and urine, blood for blood culture and seeding on to the laboratory.

6. Specific symptoms of botulism are:

- a. dry mouth;
- b. bulbar paresis (dysarthria, dysphagia, nasal regurgitation);
- c. double vision (diplopia), loss of accommodation;
- d. paralysis of facial muscles;
- e. increased salivation;
- f. paralysis of respiratory muscles.

7. The source of staphylococcal toxcosis infections can be:

- a. duck eggs;
- b. cows with mastitis;
- c. people with purulent-inflammatory diseases of the skin of the hands;
- d. people with pharyngitis;
- e. pastry.

8. Green and sprouted potatoes contain a toxic substance: e. falloidine.

- a. fagine; c. solanine:
- b. phazine; d. amygdaline;

KEY OUESTIONS

1. Food poisoning of non-microbial nature, their classification.

2. Poisoning with chemical impurities (pesticides, nitrogen-containing compounds, heavy metals). Prevention.

- 3. Food poisoning with products of plant and animal origin prevention.
- 4. Poisoning with poisonous mushrooms and plants prevention.
- 5. Tactics of the doctor in detecting food poisoning.

Notes and problems solving



Student's signature_____

Teacher's signature

HYGIENIC BASICS OF HUMAN NUTRITION IN CASE OF MAJOR SOMATIC DISEASES. HYGIENIC PRINCIPLES OF MEDICAL AND DIETARY NUTRITION

	Date
]	Learning objectives:
2	1. To learn hygienic principles of dietary and therapeutic human nutrition.
-	2. To four characteristics of the main thorapour chois.
	THEORETICAL SECTION
ve a a	lefinition
(Clinical nutrition —
]	Basics of clinical nutrition:
-	
~	1
4	1 2
4	1 2 3
	1.
2	1.
2	1.
	1.

1.	
2.	
3.	

Fill the table

Diets

Type of diet	Appointment	Characteristics

End of the table

Type of diet	Appointment	Characteristics

Each diet should include:

1.	
2.	
3.	
4.	
5.	
6.	

KEY QUESTIONS

- Hygienic principles of dietary and therapeutic human nutrition.
 Characteristics of the main therapeutic diets.
 Catering in healthcare organizations.

- 4. Features of nutrition in coronavirus infection.

Student's signature_____

Teacher's signature_____

NUTRITIONAL HYGIENE. FINAL CLASS.

CONTROL QUESTIONS

1. Nutrition hygiene: definition, purpose, objectives. The contribution of scientists in the development of the science of nutrition. Modern problems of nutrition. Types of food.

2. Rational (adequate) nutrition. The laws of rational (adequate) nutrition.

3. The law of energy adequacy nutrition. Methods for determination of energy expenditure and methods of determining the actual consumption of nutrient energy.

4. The law of nutrient adequacy of nutrition. Classification of nutrients. Formula of balanced diet.

5. The proteins and their role in human nutrition. Foods that are the main sources of complete protein.

6. Fats and carbohydrates and their role in human nutrition. Foods that are major sources of fat and carbohydrates.

7. The law of enzymatic adequacy of nutrition, the consequences of its violation. The reasons of enzymopathies occurrence.

8. The law of biotic nutrition adequacy. Food poisoning: definition, classification, characteristic.

9. Food poisoning of microbial nature: foodborne diseases, characteristic. Preventive measures.

10. Food poisoning of microbial nature: toxicosis, characteristic. Preventive measures.

11. Food poisoning of microbial nature: mycotoxicoses, ficotoxicosis, their characteristic. Preventive measures.

12. Food poisoning is no of microbial nature. Chemical toxicity (pesticides, nitrogen compounds, lead): characteristic, prevention measures.

13. Food poisoning from products of plant and animal origin. Poisoning by poisonous mushrooms its prevention.

14. Actions physician in case of revealing food poisoning.

15. The law of biorythmological adequacy of nutrition. Principles of rational diet.

16. Nutritional standards for human for energy and nutrients. The criterion for determining.

17. Nutritional status. Classification of nutritional status.

18. Stages and methods of studying the nutritional status.

19. Alimentary diseases: definition, causes, classification.

20. Diseases of protein-energy malnutrition: classification, diagnosis, clinical manifestation, prevention.

21. Inadequate (insufficient) nutritional status. Hypotrophy, alimentary dystrophy and dwarfism: clinical symptoms, prevention

22. Excess nutritional status. Obesity: causes, classification, diagnosis, prevention. Alternative and reduced diet.

23. Vitamin C deficiency. Characteristics of C and hypo- and avitaminous states, diagnosis, treatment, prevention.

24. Nutritional diseases caused by deficiency in the diet of vitamins B: diagnosis, treatment, prevention.

25. Vitamin A deficiency: diagnosis, treatment, prevention.

26. Microelementoses typical for the population of the Republic of Belarus.

27. Hyposelenosis: clinical manifestations, prevention.

28. Iodine deficiency disease: clinical manifestations, prevention.

EXAMPLES OF CONTROL CASE TASKS

1. Student (female, age — 20, body weight — 55 kg) study at the university 10 hours (duration of sleeping is 6 hours, high-intensity aerobic dancing — 1 hour, housework — 6 hours, low activity — 1 hour). Calculate total energy expenditure using physical activity levels. Intakes 80 g of proteins, 110 g of fats μ 320 g of carbohydrates. Calculate total energy expenditure, requirements for macronutrients, actual energy intake. Compare actual energy intake and total energy expenditure, actual intake of nutrients and his requirements. Make a conclusion about diet.

2. The daily diet of patient contains 160 g of cabbage. The content of vitamin C in cabbage is 50 mg per 100 g. During cooking average vitamin C losses is 55 %. There is a decrease of skin capillary resistance. Excretion of ascorbic acid in the urine was 0.4 mg/h. Assess providing the body with vitamin C.

3. During assessing the nutritional status studied physical development of students. The obtained data were compared with individual arithmetic surveyed group (female, 20 years, time of working, sleeping, social activity 8 hours each):

Sign	Individual data (a)	М	σ	Difference between individual data and M (a-M)	Sigma deviation (a-M)/σ
Height	161	165	±2,3		
Weight	51	60	±1,6		
Chest circumference	79	84	±1,9		

Draw a profile of physical development of student and assess it.

FACTORS OF THE WORKING ENVIRONMENT. DUST-RELATED OCCUPATIONAL DISEASES, THEIR PREVENTION. METHODS FOR STUDYING THE DUST CONTENT OF AIR. CHEMICAL SUBSTANCIES AS AN OCCUPATIONAL HAZARD. HYGIENIC ASSESSMENT OF THE IMPACT OF OCCUPATIONAL NOISE AND VIBRATION ON THE BODY. PREVENTION MEASURES. INFRARED RADIATION AS AN OCCUPATIONAL HAZARD. MEASURES TO PREVENT THE ADVERSE EFFECTS OF THERMAL RADIATION

Date_____

Learning objectives:

1. Studying goals and objectives of occupational hygiene.

2. Studying hygienic characteristics of working conditions in the industry.

3. Studying classification of harmful factors. Factors of the labor process.

4. Studying hygienic standards of working conditions. Occupational diseases and poisoning: types, definitions.

5. Studying hygienic characteristics of industrial dust, dust associated occupational diseases and their prevention.

6. Studying harmful effects of occupational noise and prevention of diseases related to noise.

7. Studying harmful effects of occupational vibration and prevention of diseases related to vibration.

8. Studying harmful effects of occupational heat and prevention of diseases related to occupational heat.

THEORETICAL SECTION

Give a definition

Occupational hygiene is _____

The purpose of occupational hygiene: _____

The tasks of occupational hygiene: ______

Fill the table

Classification of harmful occupational factors

Group of factors	Examples of factors

Give a definition

Workload —	
Tension of work —	
Professional disease —	
Professional poisoning —	
Acute professional poisoning —	
Chronic professional poisoning —	
Main sources of dust at work:	

Fill the tables

Dust professional pathology

Diseases	Examples
Dust respiratory diseases	
Dust diseases of the eyes	
Dust diseases of the skin	

Prevention of dust-related diseases

Aimed at technical process	Aimed at occupational environment	Aimed at employees

Main sources of noise at work: ______

Fill the tables

Noise-related health effects

Target	Effects
Ears	
Physiological effects	
Psycological effects	

Prevention of noise-related diseases

Aimed at technical process	Aimed at occupational environment	Aimed at employees

Main sources of hand-arm vibration at work: _____

Fill the tables

Vibration-related health effects

Whole-body vibration	Hand-arm vibration

Prevention of vibration-related diseases

Aimed at technical process	Aimed at occupational environment	Aimed at employees

Fill the tables

Occupational heat-related health effects

Health effects	Signs
Heat stroke	
Heat exhaustion	
Heat cramps	
Heat rash	

Aimed at technical process	Aimed at occupational environment	Aimed at employees

Prevention of heat-related diseases

KEY QUESTIONS

1. The purpose and objectives of occupational hygiene.

2. Classification of harmful factors. Factors of the labor process (severity and intensity).

3. Hygienic standards of working conditions. Occupational diseases and poisoning: types, definitions.

4. Hygienic characteristics of industrial dust.

- 5. Dust associated occupational diseases and their prevention.
- 6. Harmful effects of occupational noise and prevention of diseases related to noise.
- 7. Harmful effects of occupational vibration and prevention of diseases related to vibration.

8. Harmful effects of occupational heat and prevention of diseases related to occupational heat.

Student's signature_____

Teacher's signature_____

HYGIENIC CHARACTERISTICS OF WORKING CONDITIONS IN INDUSTRY AND AGRICULTURE. THE PROCEDURE FOR CONDUCTING PREVENTIVE MEDICAL EXAMINATIONS AT WORK. INVESTIGATION OF OCCUPATIONAL DISEASES. THE ROLE OF PREVENTIVE MEDICAL EXAMINATIONS IN THE PREVENTION OF OCCUPATIONAL DISEASES

Learning objectives:

Date_____

1. Hygienic characteristics of working conditions in the industry. Classification of harmful factors. Factors of the labor process.

2. Studying classification of medical examinations (health checks).

2. Studying purposes of medical examinations (health checks).

3. Studying steps of the investigating a worksite incident.

THEORETICAL SECTION

Fill the table

Health checks (medical examination)

Type of medical examination	Purposes of medical examination

Working conditions based on the hygienic criteria, divided into four classes:

Class 1:_	 	
Class 2:	 	
Class 3:	 	
Class 4:		

According to the degree of deviation from the parameters of the factors hygienic standards and intensity of the changes in the body of harmful working conditions of workers are divided into 4 degree of harmfulness:

Grade 1 Class 3 (3.1)	
Grade 2 Class 3 (3.2)	
Grade 3 Class 3 (3.3)	
Grade 4 Class 3 (3.4)	
Prevention of professional intoxications and diseases includes realization:	
Legislative actions	
Organizational actions	
Technological actions	
5	
Sanitary-technical actions	
reatment-preventive actions	

TRAINING TEST

1. The section of preventive medicine studying (investigating) influence on a human body of labor process and factors of the industrial environment, for the purpose of a scientific substantiation of specifications and means of preventive maintenance of occupational diseases and other adverse consequences of influences of working conditions on working —

	heept und his definition.			
1 Working conditions	A. is the factor of environment, labor process which can cause an acute			
1. working conditions	disease or sudden sharp deterioration of a state of health or even death.			
2. Dangerous	B. set of factors of labor process and the industrial environment in			
occupational factor	which the activity of the person is realized.			
3. Safe working conditions	C. conditions at which influence on working harmful and dangerous			
	production factors is excluded or their levels do not exceed hygienic			
	specifications.			
Answer:				

2. Correlate the concept and its definition:

3. Acute	profess	ional	C. is the	disease which	h has arisen	n after unitary(single) influence
poisoning			of harmf	ful substance of	on a worker	r.
Answer:						
Soloct the nk	veical har	տքսլ ք	octors.			
a tempe	ysical fiarr rature	c n	actors. athogenic	microorganis	me	e occupational noise:
h antibi	b antibiotics: d occupational heat:				,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	f vibration
Select the ch	emical har	u. 0 mful f	factors	lai neat,		1. vioration.
a antibi	otics.	mui		c heat		e enzymes
b. patho	genic micro	organ	isms:	d. industrial	noise:	e. enzymes.
. Examples of	hazardous	s dusts	s in the w	orkplace incl	ude:	
a. mold	fungi and s	pores;		c. vegetable	dusts;	
b. metal	lic dusts;	L /		d. mineral d	usts.	
. Pneumoconi	osis are oce	cupati	onal dise	ases. caused	bv:	
a. heat;		1	d. noi	se;	J	
b. vibrat	ion;		e. enz	zymes;		
c. dusts;			f. patl	hogenic micro	organisms.	
3. Prevention of	f occupatio	onal d	ust-relate	ed diseases in	cludes:	
a. using	sound-abso	orbent	coverings			
b. perso	nal protectiv	ve equ	ipment;			
c. provid	ling guidan	ce on a	seat main	tenance;		
d. trainir	ng workers	in goo	d work p	ractices;		
e. local e	exhaust ven	tilatio	n;			
f. desigr	to minimiz	ze who	ole-body v	vibration;		
g. reduc	ing airborne	e noise	e;			
h. elimiı	nation at the	e sourc	e.			
). Health effect	s of whole	-body	vibratio	n:		
a. back o	lisorders;		c. aco	oustic trauma;		
b. heada	che;		d. los	s of balance.		
0. Health effe	cts of hand	-arm	vibration	1:		
a. tingliı	ng and loss	of sen	sation in t	the fingers;		
b. shift i	n the thresh	nold of	hearing;			
c. bone	cysts in fing	gers an	d wrists;			
d. loss o	f grip streng	gth.				
1. Prevention	of occupat	ional	vibration	-related disea	ases:	
a. prohil	vition of over	ertime	work;			
b. meası	re vibration	n expo	sure;			
c. pre-er	nployment	screen	ing;			
d use of	power cha	incour				

concentration or doses of harmful substance.

B. the disease caused by the influence of harmful working

3. Correlate the concept and its definition: A. the disease developing after regular, long influence of small

conditions.

professional

1. Professional disease

2. Chronic poisoning

12. Heat-related illnesses are:

a. stroke;

c. cramps;

e. rash f. back disorders

b. pneumoconiosis; d. exhaustion

13. The main health effects of occupational noise include:

- a. bone cysts in fingers and wrists;
- b. acoustic trauma;
- c. noise-induced hearing loss;
- d. production of hormones that affect the cardiovascular system;
- e. shift in the threshold of hearing;
- f. measure vibration exposure.

14. Prevention of occupational noise-related diseases:

a. reducing airborne noise by means such as shields, enclosures, and sound-absorbent coverings;

- b. appropriate work schedules with adequate rest periods;
- c. local exhaust ventilation;
- d. individual protection measures;
- e. reducing dust concentration;
- f. elimination at the source.

KEY QUESTIONS

- 1. Harmful occupational factors in the work in industry and agriculture: impact on health.
- 2. Harmful occupational factors in the work in industry and agriculture: prevention.
- 3. Hygienic standards of working conditions.
- 4. Medical examinations of employees: types and terms of their conduct.
- 5. Role of preventive medical examinations in the prevention of occupational diseases

Student's signature_____

Teacher's signature_____

FINAL CLASS ON TOPICS «NUTRITIONAL HYGIENE» AND «OCCUPATIONAL HYGIENE»

NUTRITIONAL HYGIENE. CONTROL QUESTIONS

1. Nutrition hygiene: definition, purpose, objectives. The contribution of scientists in the development of the science of nutrition. Modern problems of nutrition. Types of food.

2. Rational (adequate) nutrition. The laws of rational (adequate) nutrition.

3. The law of energy adequacy nutrition. Methods for determination of energy expenditure and methods of determining the actual consumption of nutrient energy.

4. The law of nutrient adequacy of nutrition. Classification of nutrients. Formula of balanced diet.

5. The proteins and their role in human nutrition. Foods that are the main sources of complete protein.

6. Fats and carbohydrates and their role in human nutrition. Foods that are major sources of fat and carbohydrates.

7. The law of enzymatic adequacy of nutrition, the consequences of its violation. The reasons of enzymopathies occurrence.

8. The law of biotic nutrition adequacy. Food poisoning: definition, classification, characteristic.

9. Food poisoning of microbial nature: foodborne diseases, characteristic. Preventive measures.

10. Food poisoning of microbial nature: toxicosis, characteristic. Preventive measures.

11. Food poisoning of microbial nature: mycotoxicoses, ficotoxicosis, their characteristic. Preventive measures.

12. Food poisoning is no of microbial nature. Chemical toxicity (pesticides, nitrogen compounds, lead): characteristic, prevention measures.

13. Food poisoning from products of plant and animal origin. Poisoning by poisonous mushrooms its prevention.

14. Actions physician in case of revealing food poisoning.

15. The law of biorythmological adequacy of nutrition. Principles of rational diet.

16. Nutritional standards for human for energy and nutrients. The criterion for determining.

17. Nutritional status. Classification of nutritional status.

18. Stages and methods of studying the nutritional status.

19. Alimentary diseases: definition, causes, classification.

20. Diseases of protein-energy malnutrition: classification, diagnosis, clinical manifestation, prevention.

21. Inadequate (insufficient) nutritional status. Hypotrophy, alimentary dystrophy and dwarfism: clinical symptoms, prevention

22. Excess nutritional status. Obesity: causes, classification, diagnosis, prevention. Alternative and reduced diet.

23. Vitamin C deficiency. Characteristics of C and hypo- and avitaminous states, diagnosis, treatment, prevention.

24. Nutritional diseases caused by deficiency in the diet of vitamins B: diagnosis, treatment, prevention.

25. Vitamin A deficiency: diagnosis, treatment, prevention.

26. Microelementoses typical for the population of the Republic of Belarus.

27. Hyposelenosis: clinical manifestations, prevention.

28. Iodine deficiency disease: clinical manifestations, prevention.

OCCUPATIONAL HYGIENE. CONTROL QUESTIONS

- 1. The purpose, tasks and objectives of occupational hygiene.
- 2. Hygienic characteristics of working conditions in the industry.
- 3. Classification of harmful factors. Factors of the labor process.
- 4. Hygienic standards of working conditions.
- 5. Occupational diseases and poisoning: types, definition.
- 6. Hygienic characteristics of industrial dust.
- 7. Sources of industrial dust at workplace.
- 8. Dust associated occupational diseases and their prevention. Pneumoconiosis.
- 9. Prevention of harmful effects of dust at workplace.
- 10. Noise as harmful industrial factors.
- 11. Sources of noise at workplace.
- 12. Physiological and psychological effects of noise.
- 13. Prevention of harmful effects of noise.
- 14. Vibration as harmful industrial factor.
- 15. Sources of hand-arm and whole-body vibration at workplace.
- 16. Health effects of vibration.
- 17. Prevention of harmful effects of hand-arm and whole-body vibration at workplace.
- 18. Hygienic characteristic of chemical factors. The routes of entry of chemical substances.
- 19. Health effects of chemical factors.
- 20. Prevention of occupational poisonings and hazard effects of chemical substances.
- 21. Hygienic characteristic of infrared radiation.
- 22. Health effects of heat exposure. Types of heat-related illnesses.
- 22. Prevention of infrared radiation associated occupational diseases.
- 23. Types and purposes of medical examinations (health checks).

CONTENT

Physical properties of atmospheric air. Research methods and hygienic assessment of microclimate parameters in residential premises and medical institutions (temperature, humidity)
Physical properties of atmospheric air. Research methods and hygienic assessment of microclimate parameters in residential premises and medical institutions (air velocity, atmospheric pressure)
Physical properties of atmospheric air. Hygienic assessment of the combined action of meteorological factors on the human body
Physical properties and chemical composition of atmospheric air. Final class
Hygienic assessment of the impact of placement conditions on human health. Research methods and hygienic assessment of natural lighting in enclosed premises
Hygienic assessment of the impact of placement conditions on human health. Research methods and hygienic assessment of artificial lighting in enclosed premises
Hygienic assessment of the impact of placement conditions on human health. Natural and artificial ventilation, types and their hygienic characteristics. Research methods and hygienic assessment of carbon dioxide concentration. Hygienic assessment of ventilation indicators: air cube, ventilation volume, air exchange rate.
Hygienic assessment of the impact of placement on human health Final class 28
Hygienic assessment of drinking water quality. Research methods and hygienic assessment
of physical, organoleptic and chemical indicators of water quality
Hygienic assessment of drinking water quality. Methods for improving water quality
Hygienic assessment of drinking water quality. Final class
Hygienic assessment of soil
Hygiene of healthcare organizations. Hygienic assessment of the general plan of the healthcare organization
Hygiene of healthcare organizations. Hygienic requirements for the device and equipment of the ward section of the healthcare organization department
Hygiene of children and adolescents. Assessment of physical development of children and adolescents
Final class on topics «The structure of the human environment and it's hygienic significance», «Hygiene of healthcare organizations», «Hygiene of children and adolescents»
Hygienic assessment of energy value and nutritional adequacy of the diet. Criteria for rationing the body's needs for nutrition, physiological requirements for the nutrition of the population. Methods for determining the body's needs for energy and nutrients
Hygienic assessment of the energy value and nutritional adequacy of the diet. Calculation of the actual nutrient intake and energy value of the diet according to the 24-hour re-call method 58
Hygienic assessment of the energy value and nutritional adequacy of the diet. Assessment of the adequacy of actual diet to the needs of the human body, recommendations for rationalizing nutrition
Hygienic assessment of nutritional status. Determination of somatometric, somatoscopic, physiometric indicators

Hygienic assessment of nutritional status. Recommendations for optimizing nutritional status
Hygienic assessment of the provision of the human body with vitamins. Recommendations for optimizing the vitamin supply of the human body
Prevention of food poisoning of microbial etiology
Prevention of non-microbial food poisoning
Hygienic basics of human nutrition in case of major somatic diseases. Hygienic principles of medical and dietary nutrition
Nutritional hygiene. Final class
Factors of the working environment. Dust-related occupational diseases, their prevention. Methods for studying the dust content of air. Chemical substancies as an occupational hazard. Hygienic assessment of the impact of occupational noise and vibration on the body. Prevention measures. Infrared radiation as an occupational hazard. Measures to prevent the adverse effects of thermal radiation
Hygienic characteristics of working conditions in industry and agriculture. The procedure for conducting preventive medical examinations at work. Investigation of occupational diseases. The role of preventive medical examinations in the prevention of occupational diseases
Final class on topics «Nutritional hygiene» and «Occupational hygiene»